arXiv:0806.3615v2 [math.RT] 4 Aug 2008

A QUIVER CONSTRUCTION OF SYMMETRIC CRYSTALS

NAOYA ENOMOTO

Abstract. In the papers [EK1], [EK2] and [EK3] with Masaki Kashiwara, the author introduced the notion of symmetric crystals and presented the Lascoux-Leclerc-Thibon-Ariki type conjectures for the affine Hecke algebras of type B. Namely, we conjectured that certain composition multiplicities and branching rules for the affine Hecke algebras of type B are described by using the lower global basis of symmetric crystals of $V_\theta(\lambda)$. In the present paper, we prove the existence of crystal bases and global bases of $V_\theta(0)$ for any symmetric quantized Kac-Moody algebra by using a geometry of quivers (with a Dynkin diagram involution). This is analogous to George Lusztig's geometric construction of U_v^- and its lower global basis.

1. Introduction

1.1. Let K_n^{AHA} be the Grothendieck group of the affine Hecke algebra $H_n(q)$ of type A_n and set $K^{\mathrm{AHA}} = \bigoplus_{n\geq 0} K_n^{\mathrm{AHA}}$. Generalizing the LLT conjecture [LLT] for the Hecke algebra of type A, S. Ariki [Ari] proved that $K^{\mathrm{AHA}} \otimes_{\mathbb{Z}} \mathbb{C}$ is isomorphic to $U^-(\mathfrak{g})$ as $U^-(\mathfrak{g})$-modules. Here $\mathfrak{g} = \widehat{\mathfrak{sl}}_{\ell-1}$ or $\mathfrak{gl}_\infty$ according that the parameter q of the affine Hecke algebras of type A is a primitive ℓ-th root of unity or not a root of unity. This isomorphism sends the irreducible modules of the affine Hecke algebras to the specialization of the upper global basis of $U_v^-(\mathfrak{g})$ at $v=1$. His proof is based on two results in the geometric representation theory. One is the equivariant K-theoretic description of the irreducible and standard modules of the affine Hecke algebras by Chriss-Ginzburg and Kazhdan-Lusztig, and the other is G. Lusztig's geometric construction [Lus1] of the lower global basis of $U_v^-(\mathfrak{g})$. Lusztig's theory is summarized as follows.

Let $\mathfrak{g}$ be a symmetric Kac-Moody algebra and I an index set of simple roots of $\mathfrak{g}$. For a fixed set of arrows Ω, we consider (I,Ω) as a (finite) oriented graph. We call (I,Ω) a quiver. For an I-graded vector space $\mathbf{V}$, we define the moduli space of representations of quiver (I,Ω) by

$$\mathbf{E}_{\mathbf{V},\Omega} = \bigoplus_{i \xrightarrow{\Omega} j} \mathrm{Hom}(\mathbf{V}_i, \mathbf{V}_j).$$

The algebraic group $G_{\mathbf{V}} = \prod_{i\in I} GL(\mathbf{V}_i)$ acts on $\mathbf{E}_{\mathbf{V},\Omega}$. Lusztig introduced a certain full subcategory $\mathscr{Q}_{\mathbf{V},\Omega}$ of $\mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$ where $\mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$ is the bounded derived category of constructible complexes of sheaves on $\mathbf{E}_{\mathbf{V},\Omega}$ (for the definition, see section 3). Let $K(\mathscr{Q}_{\mathbf{V},\Omega})$ be the Grothendieck group of $\mathscr{Q}_{\mathbf{V},\Omega}$. He constructed the induction operators f_i and the restriction operators e'_i on the Grothendieck group $K_\Omega := \oplus_{\mathbf{V}} K(\mathscr{Q}_{\mathbf{V},\Omega})$, where $\mathbf{V}$ runs over the isomorphism classes of I-graded vector spaces. He proved the following theorem.

Theorem 1.1 (Lusztig).

(i) The operators e'_i and f_i define the action of the reduced v-analogue $B_v(\mathfrak{g})$ of $\mathfrak{g}$ on $K_\Omega \otimes_{\mathbb{Z}[v,v^{-1}]} \mathbb{Q}(v)$, and $K_\Omega \otimes_{\mathbb{Z}[v,v^{-1}]} \mathbb{Q}(v)$ is isomorphic to $U_v^-(\mathfrak{g})$ as a $B_v(\mathfrak{g})$-module.

The author is partially supported by JSPS Research Fellowships for Young Scientists.

The involution induced by the Verdier duality functor coincides with the bar involution on $U_v^-(\mathfrak{g})$.

(ii) The simple perverse sheaves in $\bigoplus_{\mathbf{V}} \mathscr{Q}_{\mathbf{V},\Omega}$ *give the lower global basis of* $U_v^-(\mathfrak{g})$.

1.2. Recently in [EK1] and [EK2] with M. Kashiwara, the author presented an analogue of the LLTA conjecture for the affine Hecke algebra of *type B*. In [EK2], we considered $U_v(\mathfrak{g})$ and its Dynkin diagram involution θ and constructed an analogue $B_\theta(\mathfrak{g})$ of the reduced v-analogue $B_v(\mathfrak{g})$ (for the definition, see Definition 2.9 below). We gave a $B_\theta(\mathfrak{g})$-module $V_\theta(\lambda)$ for a dominant integral weight λ such that $\theta(\lambda) = \lambda$, which is an analogue of the $B_v(\mathfrak{g})$-module $U_v^-(\mathfrak{g})$ (for the definition, see Definition 2.10 below). We defined the notion of symmetric crystals and conjectured the existence of the global basis. In the case $\mathfrak{g} = \mathfrak{gl}_\infty$, $I = \mathbb{Z}_{\mathrm{odd}}$, $\theta(i) = -i$ and $\lambda = 0$, we constructed the PBW type basis and the lower (and upper) global basis parametrized by the θ-restricted multi-segments. We conjectured that irreducible modules of the affine Hecke algebras of type B are described by the global basis associated to the symmetric crystals.

1.3. In this paper, we construct the lower global basis for the symmetric crystals by using a geometry of quivers (with a Dynkin diagram involution). Hence for any symmetric quantized Kac-Moody algebra $U_v(\mathfrak{g})$, we establish the existence of a crystal basis and a global basis for $V_\theta(0)$.

We introduce the notion of θ-quivers. This is a quiver (I, Ω) with an involution $\theta : I \to I$ (and $\theta : \Omega \to \Omega$) satisfing some conditions (see Definition 4.1). This notion is partially motivated by Syu Kato's construction [Kt] of the irreducible representations of the affine Hecke algebras of type B.

We also introduce the θ-symmetric I-graded vector spaces. This is an I-graded vector space $\mathbf{V} = (\mathbf{V}_i)_{i \in I}$ endowed with a non-degenerate symmetric bilinear form such that $\mathbf{V}_i$ and $\mathbf{V}_j$ are orthogonal if $j \neq \theta(i)$. For a θ-quiver (I, Ω) and a θ-symmetric I-graded vector space $\mathbf{V}$, we define the moduli space ${}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ of representations of (I, Ω) adding a skew-symmetric condition on $\mathbf{E}_{\mathbf{V},\Omega}$ with respect to the involution θ.

Similarly to Lusztig's arguments, we consider a certain full subcategory ${}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$ of $\mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$ and its Grothendieck group ${}^\theta K_{\mathbf{V},\Omega}$. We define the induction operators F_i and the restriction operators E_i on ${}^\theta K_\Omega := \oplus_{\mathbf{V}} {}^\theta K_{\mathbf{V},\Omega}$ where $\mathbf{V}$ runs over the isomorphism classes of the θ-symmetric I-graded vector spaces. We prove the following main theorem which is an analogous result of Theorem 1.1.

Theorem 1.2 (Theorem 5.12)**.** ${}^\theta K_\Omega \otimes_{\mathbb{Z}[v,v^{-1}]} \mathbb{Q}(v) \cong V_\theta(0)$ *as* $B_\theta(\mathfrak{g})$*-modules. The simple perverse sheaves in* ${}^\theta K_\Omega$ *give a lower global basis of* $V_\theta(0)$.

Though Lusztig proved Theorem 1.1 using some inner product on K_Ω, we prove Theorem 1.2 using a criterion of crystals (Theorem 2.14) and certain estimates for the actions of E_i and F_i on simple perverse sheaves (Theorem 5.3).

This paper is organized as follows.

In section 2, we recall some results on the quantum enveloping algebras, the theory of the crystal bases and the global bases, the notion of symmetric crystals and known results of perverse sheaves and Fourier-Sato-Deligne transforms. Especially, we recall an important criterion of crystals in Theorem 2.14. We use this in our proof of existence of the crystal basis of $V_\theta(0)$.

In section 3, we give a quick review on Lusztig's construction of $U_v^-(\mathfrak{g})$ and its lower global basis.

In section 4, we introduce the notion of θ-quivers and θ-symmetric I-graded vector spaces. We define the category ${}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$ and the induction operators F_i and the restriction operators

E_i. We calculate actions of E_i and F_i on ${}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega}$. We also prove that E_i and F_i commute with the Fourier-Sato-Deligne transforms.
In section 5, we introduce the Grothendieck group ${}^{\theta}K_{\Omega}$ and show three key results. First, we calculate the commutation relations of E_i and F_i. Second, we give certain estimates of coefficients with respect to the action of E_i and F_i on simple perverse sheaves. These estimates satisfy the condition in Theorem 2.14. Third, we prove the invariance of simple perverse sheaves with respect to the Verdier duality functor. Combining these results we prove the main theorem.

Remark 1.3. We give two remarks on a difference from the "folding" procedure and an overlap with perverse sheaves arising from graded Lie algebras by Lusztig.

(i) Our construction is completely different from Lusztig's construction, "Quiver with automorphisms", in his book [Lus3, Chapter.12-14].
He considered actions $a : I \to I$ and $a : H \to H$ induced from a finite cyclic group C generated by a. Put an orientation Ω such that $\mathrm{out}(a(h)) = a(\mathrm{out}(h))$ and $\mathrm{in}(a(h)) = a(\mathrm{in}(h))$. He said this orientation "compatible". Let $\mathcal{V}^a$ be the category of I-graded vector spaces $\mathbf{V}$ such that $\dim \mathbf{V}_i = \dim \mathbf{V}_{a(i)}$ for any $i \in I$. For $\mathbf{V} \in \mathcal{V}^a$, a induces a natural automorphism on $\mathbf{E}_{\mathbf{V},\Omega}$ and a functor $a^* : \mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega}) \to \mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$. He introduced "$C$-equivariant" simple perverse sheaves (B, ϕ), where B is a perverse sheaf on $\mathbf{E}_{\mathbf{V},\Omega}$ and $\phi : a^*B \cong B$. Then he proved that the set $\sqcup_{\mathbf{V}\in\mathcal{V}^a}\mathbf{B}_{\mathbf{V},\Omega}$ of C-equivariant perverse sheaves gives a lower global basis of $\mathbf{U}_v^-(\mathfrak{g})$. Here $\mathfrak{g}$ has a non-symmetric Cartan matrix which is obtained by the "folding" procedure with respect to the C-action on I.
But in our construction, a θ-orientation is not a compatible orientation. Moreover the most essential difference is that his construction has no skew-symmetric condition in our sence. Hence the set of simple perverse sheaves ${}^{\theta}\mathscr{P}_{\mathbf{V},\Omega}$ and the space ${}^{\theta}K_{\Omega} \otimes_{\mathbb{Z}[v,v^{-1}]} \mathbb{Q}(v) \cong V_{\theta}(0)$ are different from $\mathbf{B}_{\mathbf{V},\Omega}$ and $\mathbf{U}_v^-(\mathfrak{g})$, respectively. The explicit crystal structure of $V_{\theta}(0)$ is unknown except for the case $\mathfrak{g} = \mathfrak{gl}_{\infty}$, $I = \mathbb{Z}_{\mathrm{odd}}$ and $\theta(i) = -i$ in [EK2].

(ii) In some special case, the lower global basis constructed in this paper is obtained by Lusztig ([Lus4] and [Lus5]). Let us consider the case $G = SO(2n, \mathbb{C})$. Let $\mathfrak{g}$ be the Lie algebra of G and T a fixed maximal torus of G. Set ε_{2i-1} $(1 \leq i \leq n)$ the fundamental characters of T. Asuume $q \in \mathbb{C}^*$ is not a root of unity. We choose a semisimple element $s \in T$ such that $\varepsilon_{2i-1}(s) \in q^{\mathbb{Z}_{\mathrm{odd},\geq 0}}$ for any i and put $d_{2i-1} = \{j | \varepsilon_{2j-1}(s) = q^{2i-1}\}$. Then the centralizer $G(s)$ of s acts on

$$\mathfrak{g}_2 := \{X \in \mathfrak{g} \mid sXs^{-1} = q^2X\}$$

which has finitely many $G(s)$-oribits. Lusztig considered the category $\mathscr{Q}(\mathfrak{g}_2)$ of semisimple $G(s)$-equivariant complex on $\mathfrak{g}_2$ and constructed the canonical basis $\mathbf{B}(\mathfrak{g}_2)$ of $K(\mathfrak{g}_2)$ which is the Grothendieck group of $\mathscr{Q}(\mathfrak{g}_2)$.
On the other hand, let us consider the θ-symmetric vector space $\mathbf{V}$ such that $\mathrm{wt}(\mathbf{V}) = \sum_{i=1}^{n} d_{2i-1}(\alpha_{2i-1}+\alpha_{-2i+1})$ and the following θ-quiver of type A_{2n} and the θ-orientation Ω:



In this case, we have $G(s) = \prod_{i=1}^{n} GL(d_{2i-1}) = {}^{\theta}\mathbf{G}_{\mathbf{V}}$ and $\mathfrak{g}_2 \cong {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}$. Thus the set ${}^{\theta}\mathscr{P}_{\mathbf{V},\Omega}$ of simple perverse sheaves conincide with $\mathbf{B}(\mathfrak{g}_2)$.

Acknowledgements. I would like to thank Masaki Kashiwara, Georg Lusztig, Susumu Ariki, Syu Kato and Yuichiro Hoshi. Masaki Kashiwara guided me to Lusztig's geometric theory and gave me many advises, comments and patient encouragement. George Lusztig pointed out the relations between his work and this paper. Syu Kato also commented on this paper and guided me to his geometric representation theory of AHA of type C_n and certain quiver presentations. Susumu Ariki gave me some comments and encouragement. Yuichiro Hoshi taught me basic concepts, examples and some techniques in algebraic geometry and derived categories till midnights at RIMS.

2. Preliminaries

2.1. **Quantum enveloping algebras.**

2.1.1. *Quantum enveloping algebras and reduced v-analogue.* We shall recall the quantized universal enveloping algebra $U_v(\mathfrak{g})$. In this paper, we treat only the symmetric Cartan matrix case. Let I be an index set (for simple roots), and Q the free $\mathbb{Z}$-module with a basis $\{\alpha_i\}_{i\in I}$. Let $(\bullet, \bullet) : Q \times Q \to \mathbb{Z}$ be a symmetric bilinear form such that $(\alpha_i, \alpha_i) = 2$ and $(\alpha_i, \alpha_j) \in \mathbb{Z}_{\le 0}$ for $i \neq j$. Let v be an indeterminate and set $\mathbf{K} := \mathbb{Q}(v)$. We define its subrings $\mathbf{A}_0$, $\mathbf{A}_\infty$ and $\mathbf{A}$ as follows.

$$\begin{aligned}
\mathbf{A}_0 &= \{f \in \mathbf{K} \mid f \text{ is regular at } v = 0\}, \\
\mathbf{A}_\infty &= \{f \in \mathbf{K} \mid f \text{ is regular at } v = \infty\}, \\
\mathbf{A} &= \mathbb{Q}[v, v^{-1}].
\end{aligned}$$

Definition 2.1. *The quantized universal enveloping algebra $U_v(\mathfrak{g})$ is the $\mathbf{K}$-algebra generated by elements e_i, f_i and invertible elements t_i $(i \in I)$ with the following defining relations.*

(1) *The t_i's commute with each other.*

(2) $t_j e_i t_j^{-1} = v^{(\alpha_j,\alpha_i)} e_i$ *and* $t_j f_i t_j^{-1} = v^{-(\alpha_j,\alpha_i)} f_i$ *for any $i, j \in I$.*

(3) $[e_i, f_j] = \delta_{ij} \dfrac{t_i - t_i^{-1}}{v - v^{-1}}$ *for $i, j \in I$.*

(4) (v-Serre relation) *For $i \neq j$,*

$$\sum_{k=0}^{b} (-1)^k e_i^{(k)} e_j e_i^{(b-k)} = 0, \ \sum_{k=0}^{b} (-1)^k f_i^{(k)} f_j f_i^{(b-k)} = 0.$$

Here $b = 1 - (\alpha_i, \alpha_j)$ and

$$e_i^{(k)} = e_i^k/[k]_v!\,,\ f_i^{(k)} = f_i^k/[k]_v!\ ,\ [k]_v = (v^k - v^{-k})/(v - v^{-1})\,,\ [k]_v! = [1]_v \cdots [k]_v\,.$$

Let us denote by $U_v^-(\mathfrak{g})$ the subalgebra of $U_v(\mathfrak{g})$ generated by the f_i's.

Let e_i' and e_i^* be the operators on $U_v^-(\mathfrak{g})$ defined by

$$[e_i, a] = \frac{(e_i^* a) t_i - t_i^{-1} e_i' a}{v - v^{-1}} \quad (a \in U_v^-(\mathfrak{g})).$$

These operators satisfy the following formulas similar to derivations:

$$e_i'(ab) = (e_i' a) b + (\mathrm{Ad}(t_i) a) e_i' b.$$

The algebra $U_v^-(\mathfrak{g})$ has a unique symmetric bilinear form $(\bullet, \bullet)$ such that $(1, 1) = 1$ and

$$(e_i' a, b) = (a, f_i b) \quad \text{for any } a, b \in U_v^-(\mathfrak{g}).$$

It is non-degenerate. The left multiplication operator f_j and e'_i satisfy the commutation relations

$$e'_i f_j = v^{-(\alpha_i,\alpha_j)} f_j e'_i + \delta_{ij},\ e^*_i f_j = f_j e^*_i + \delta_{ij}\operatorname{Ad}(t_i),$$

and the e'_i's satisfy the v-Serre relations.

Definition 2.2. *The reduced v-analogue $B_v(\mathfrak{g})$ of $\mathfrak{g}$ is the $\mathbb{Q}(v)$-algebra generated by e'_i and f_i.*

2.1.2. *Review on crystal bases and global bases of U_v^-.* Since e'_i and f_i satisfy the v-boson relation, any element $a \in U_v^-(\mathfrak{g})$ can be uniquely written as

$$a = \sum_{n\geq 0} f_i^{(n)} a_n \quad \text{with } e'_i a_n = 0.$$

Here $f_i^{(n)} = \dfrac{f_i^n}{[n]_v!}$.

Definition 2.3. *We define the modified root operators $\widetilde{e}_i$ and $\widetilde{f}_i$ on $U_v^-(\mathfrak{g})$ by*

$$\widetilde{e}_i a = \sum_{n\geq 1} f_i^{(n-1)} a_n, \quad \widetilde{f}_i a = \sum_{n\geq 0} f_i^{(n+1)} a_n.$$

Theorem 2.4 ([Ka1])**.** *We define*

$$\begin{aligned} L(\infty) &= \sum_{\ell\geq 0,\, i_1,\ldots,i_\ell\in I} \mathbf{A}_0 \widetilde{f}_{i_1}\cdots\widetilde{f}_{i_\ell}\cdot 1 \subset U_v^-(\mathfrak{g}), \\ B(\infty) &= \left\{ \tilde{f}_{i_1}\cdots\tilde{f}_{i_\ell}\cdot 1 \mod vL(\infty) \mid \ell\geq 0, i_1,\cdots,i_\ell\in I \right\} \subset L(\infty)/vL(\infty). \end{aligned}$$

Then we have

(1) $\widetilde{e}_i L(\infty)\subset L(\infty)$ and $\widetilde{f}_i L(\infty)\subset L(\infty)$,
(2) $B(\infty)$ is a basis of $L(\infty)/vL(\infty)$,
(3) $\widetilde{f}_i B(\infty)\subset B(\infty)$ and $\widetilde{e}_i B(\infty)\subset B(\infty)\cup\{0\}$.

We call $(L(\infty), B(\infty))$ the crystal basis of $U_v^-(\mathfrak{g})$.

Definition 2.5. *We define $\varepsilon_i(b) := \max\{m\in\mathbb{Z}_{\geq 0} | \widetilde{e}_i^m b\neq 0\}$ for $i\in I$ and $b\in B(\infty)$.*

Let $-$ be the automorphism of $\mathbf{K}$ sending v to v^{-1}. Then $\overline{\mathbf{A}_0}$ coincides with $\mathbf{A}_\infty$.

Let V be a vector space over $\mathbf{K}$, L_0 an $\mathbf{A}$-submodule of V, L_∞ an $\mathbf{A}_\infty$- submodule, and $V_{\mathbf{A}}$ an $\mathbf{A}$-submodule. Set $E := L_0\cap L_\infty\cap V_{\mathbf{A}}$.

Definition 2.6 ([Ka1])**.** *We say that $(L_0, L_\infty, V_{\mathbf{A}})$ is* balanced *if each of L_0, L_∞ and $V_{\mathbf{A}}$ generates V as a $\mathbf{K}$-vector space, and if one of the following equivalent conditions is satisfied.*

(1) $E\to L_0/vL_0$ is an isomorphism,
(2) $E\to L_\infty/v^{-1}L_\infty$ is an isomorphism,
(3) $(L_0\cap V_{\mathbf{A}})\oplus(v^{-1}L_\infty\cap V_{\mathbf{A}})\to V_{\mathbf{A}}$ is an isomorphism.
(4) $\mathbf{A}_0\otimes_{\mathbb{Q}}E\to L_0$, $\mathbf{A}_\infty\otimes_{\mathbb{Q}}E\to L_\infty$, $\mathbf{A}\otimes_{\mathbb{Q}}E\to V_{\mathbf{A}}$ and $\mathbf{K}\otimes_{\mathbb{Q}}E\to V$ are isomorphisms.

Let $-$ be the ring automorphism of $U_v(\mathfrak{g})$ sending v, t_i, e_i, f_i to v^{-1}, t_i^{-1}, e_i, f_i.

Let $U_v(\mathfrak{g})_{\mathbf{A}}$ be the $\mathbf{A}$-subalgebra of $U_v(\mathfrak{g})$ generated by $e_i^{(n)}$, $f_i^{(n)}$ and t_i. Similarly we define $U_v^-(\mathfrak{g})_{\mathbf{A}}$.

Theorem 2.7. *$(L(\infty), L(\infty)^-, U_v^-(\mathfrak{g})_{\mathbf{A}})$ is balanced.*

Let

$$G^{\mathrm{low}}\colon L(\infty)/vL(\infty)\xrightarrow{\sim}E:=L(\infty)\cap L(\infty)^{-}\cap U_v^{-}(\mathfrak{g})_{\mathbf{A}}$$

be the inverse of $E\xrightarrow{\sim}L(\infty)/vL(\infty)$. Then $\{G^{\mathrm{low}}(b)\mid b\in B(\infty)\}$ forms a basis of $U_v^{-}(\mathfrak{g})$. We call it a (lower) *global basis*. It is first introduced by G. Lusztig ([Lus1]) under the name of "canonical basis" for the A, D, E cases.

Definition 2.8. *Let*

$$\{G^{\mathrm{up}}(b)\mid b\in B(\infty)\}$$

be the dual basis of $\{G^{low}(b)\mid b\in B(\infty)\}$ *with respect to the inner product* $(\bullet,\bullet)$. *We call it the upper global basis of* $U_v^{-}(\mathfrak{g})$.

2.2. **Symmetric Crystals.** Let θ be an automorphism of I such that $\theta^2=\mathrm{id}$ and $(\alpha_{\theta(i)},\alpha_{\theta(j)})=(\alpha_i,\alpha_j)$. Hence it extends to an automorphism of the root lattice Q by $\theta(\alpha_i)=\alpha_{\theta(i)}$, and induces an automorphism of $U_v(\mathfrak{g})$.

Definition 2.9. *Let* $B_\theta(\mathfrak{g})$ *be the* $\mathbf{K}$*-algebra generated by* E_i, F_i, *and invertible elements* $T_i (i\in I)$ *satisfying the following defining relations:*

- (i) *the* T_i*'s commute with each other,*
- (ii) $T_{\theta(i)}=T_i$ *for any* i,
- (iii) $T_iE_jT_i^{-1}=v^{(\alpha_i+\alpha_{\theta(i)},\alpha_j)}E_j$ *and* $T_iF_jT_i^{-1}=v^{(\alpha_i+\alpha_{\theta(i)},-\alpha_j)}F_j$ *for* $i,j\in I$,
- (iv) $E_iF_j=v^{-(\alpha_i,\alpha_j)}F_jE_i+(\delta_{i,j}+\delta_{\theta(i),j}T_i)$ *for* $i,j\in I$,
- (v) *the* E_i*'s and the* F_i*'s satisfy the* v*-Serre relations.*

We set $F_i^{(n)}=F_i^n/[n]_v!$.

Proposition 2.10 ([EK2, Proposition2.11.]). *Let*

$$\lambda\in P_+:=\{\lambda\in\mathrm{Hom}(Q,\mathbb{Q})\mid\lambda(\alpha_i)\in\mathbb{Z}_{\geq0}\text{ for any }i\in I\}$$

be a dominant integral weight such that $\theta(\lambda)=\lambda$.

- (i) *There exists a* $B_\theta(\mathfrak{g})$*-module* $V_\theta(\lambda)$ *generated by a non-zero vector* ϕ_λ *such that*
  - (a) $E_i\phi_\lambda=0$ *for any* $i\in I$,
  - (b) $T_i\phi_\lambda=v^{(\alpha_i,\lambda)}\phi_\lambda$ *for any* $i\in I$,
  - (c) $\{u\in V_\theta(\lambda)\mid E_iu=0\text{ for any }i\in I\}=\mathbf{K}\phi_\lambda$.

  Moreover such a $V_\theta(\lambda)$ *is irreducible and unique up to an isomorphism.*
- (ii) *There exists a unique non-degenerate symmetric bilinear form* $(\bullet,\bullet)$ *on* $V_\theta(\lambda)$ *such that* $(\phi_\lambda,\phi_\lambda)=1$ *and* $(E_iu,v)=(u,F_iv)$ *for any* $i\in I$ *and* $u,v\in V_\theta(\lambda)$.
- (iii) *There exists an endomorphism* $-$ *of* $V_\theta(\lambda)$ *such that* $\overline{\phi_\lambda}=\phi_\lambda$ *and* $\overline{av}=\bar{a}\bar{v}$, $\overline{F_iv}=F_i\bar{v}$ *for any* $a\in\mathbf{K}$ *and* $v\in V_\theta(\lambda)$.

Hereafter we assume further that

there is no $i\in I$ *such that* $\theta(i)=i$.

In [EK2], we conjectured that $V_\theta(\lambda)$ has a crystal basis. This means the following. Since E_i and F_i satisfy the v-boson relation $E_iF_i=v^{-(\alpha_i,\alpha_i)}F_iE_i+1$, we define the modified root operators:

$$\tilde{E}_i(u)=\sum_{n\geq1}F_i^{(n-1)}u_n\text{ and }\tilde{F}_i(u)=\sum_{n\geq0}F_i^{(n+1)}u_n,$$

when writing $u=\sum_{n\geq0}F_i^{(n)}u_n$ with $E_iu_n=0$. Let $L_\theta(\lambda)$ be the $\mathbf{A}_0$-submodule of $V_\theta(\lambda)$ generated by $\widetilde{F}_{i_1}\cdots\widetilde{F}_{i_\ell}\phi_\lambda$ ($\ell\geq0$ and $i_1,\ldots,i_\ell\in I$), and let $B_\theta(\lambda)$ be the subset

$$\left\{\widetilde{F}_{i_1}\cdots\widetilde{F}_{i_\ell}\phi_\lambda \mod vL_\theta(\lambda)\mid\ell\geq0,i_1,\ldots,i_\ell\in I\right\}$$

of $L_\theta(\lambda)/vL_\theta(\lambda)$.

Conjecture 2.11. *Let λ be a dominant integral weight such that $\theta(\lambda) = \lambda$.*

(1) $\tilde{F}_i L_\theta(\lambda) \subset L_\theta(\lambda)$ and $\tilde{E}_i L_\theta(\lambda) \subset L_\theta(\lambda)$,
(2) $B_\theta(\lambda)$ is a basis of $L_\theta(\lambda)/vL_\theta(\lambda)$,
(3) $\tilde{F}_i B_\theta(\lambda) \subset B_\theta(\lambda)$, and $\tilde{E}_i B_\theta(\lambda) \subset B_\theta(\lambda) \sqcup \{0\}$,
(4) $\tilde{F}_i \tilde{E}_i(b) = b$ for any $b \in B_\theta(\lambda)$ such that $\tilde{E}_i b \neq 0$, and $\tilde{E}_i \tilde{F}_i(b) = b$ for any $b \in B_\theta(\lambda)$.

Moreover we conjectured that $V_\theta(\lambda)$ has a global crystal basis. Namely we have

Conjecture 2.12. $(L_\theta(\lambda), \overline{L_\theta(\lambda)}, V_\theta(\lambda)^{low}_{\mathbf{A}})$ *is balanced. Here* $V_\theta(\lambda)^{low}_{\mathbf{A}} := U_v^-(\mathfrak{g})_{\mathbf{A}}\phi_\lambda$.

Example 2.13. Suppose $\mathfrak{g} = \mathfrak{gl}_\infty$, the Dynkin diagram involution θ of I defined by $\theta(i) = -i$ for $i \in I = \mathbb{Z}_{\mathrm{odd}}$.



And assume $\lambda = 0$. In this case, we can prove

$$V_\theta(0) \cong U_v^- / \sum_{i \in I} U_v^-(f_i - f_{\theta(i)}).$$

Moreover we can construct a PBW type basis, a crystal basis and an upper and lower global basis on $V_\theta(0)$ parametrized by "the θ-restricted multisegments". For more details, see [EK2].

2.3. **Criterion for crystals.** Let $\mathbf{K}[e, f]$ be the ring generated by e and f with the defining relation $ef = v^{-2}fe + 1$. We call this algebra the v-boson algebra. Let P be a free $\mathbb{Z}$-module, and let α be a non-zero element of P. Let M be a $\mathbf{K}[e, f]$-module. Assume that M has a weight decomposition $M = \oplus_{\xi \in P} M_\xi$ and $eM_\lambda \subset M_{\lambda+\alpha}$ and $fM_\lambda \subset M_{\lambda-\alpha}$. Asuume the following finiteness conditions:

$$\text{for any } \lambda \in P, \dim M_\lambda < \infty \text{ and } M_{\lambda+n\alpha} = 0 \text{ for } n \gg 0.$$

Hence for $u \in M$, we can write $u = \sum_{n \geq 0} f^{(n)} u_n$ with $eu_n = 0$. We define endmorphisms $\widetilde{e}$ and $\widetilde{f}$ of M by

$$\widetilde{e}u = \sum_{n \geq 1} f^{(n-1)} u_n, \quad \widetilde{f}u = \sum_{n \geq 0} f^{(n+1)} u_n.$$

Let B be a crystal with weight decomposition by P in the following sense. We have $\mathrm{wt}\colon B \to P$, $\widetilde{f}\colon B \to B$, $\widetilde{e}\colon B \to B \sqcup \{0\}$ and $\varepsilon\colon B \to \mathbb{Z}_{\geq 0}$ satisfying the following properties, where $B_\lambda = \mathrm{wt}^{-1}(\lambda)$:

(i) $\widetilde{f} B_\lambda \subset B_{\lambda-\alpha}$ and $\widetilde{e} B_\lambda \subset B_{\lambda+\alpha} \sqcup \{0\}$ for any $\lambda \in P$,
(ii) $\widetilde{f}\widetilde{e}b = b$ if $\widetilde{e}b \neq 0$, and $\widetilde{e} \circ \widetilde{f} = \mathrm{id}_B$,
(iii) for any $\lambda \in P$, B_λ is a finite set and $B_{\lambda+n\alpha} = \phi$ for $n \gg 0$,
(iv) $\varepsilon(b) = \max\{n \geq 0 \mid \widetilde{e}^n b \neq 0\}$ for any $b \in B$.

Set $\mathrm{ord}(a) = \sup\{n \in \mathbb{Z} \mid a \in v^n \mathbf{A}_0\}$ for $a \in \mathbf{K}$. We understand $\mathrm{ord}(0) = \infty$.
Let $\{G(b)\}_{b \in B}$ be a system of generators of M with $G(b) \in M_{\mathrm{wt}(b)}$. Asuume that we have expressions:

$$eG(b) = \sum_{b' \in B} E_{b,b'} G(b), \quad fG(b) = \sum_{b' \in B} F_{b,b'} G(b).$$

Now consider the following conditions for these data, where $\ell = \varepsilon(b)$ and $\ell' = \varepsilon(b')$:

$$\operatorname{ord}(F_{b,b'}) \geq 1 - \ell', \tag{2.1}$$

$$\operatorname{ord}(E_{b,b'}) \geq -\ell', \tag{2.2}$$

$$F_{b,\widetilde{f}b} \in v^{-\ell}(1 + v\mathbb{A}_0), \tag{2.3}$$

$$E_{b,\widetilde{f}b} \in v^{1-\ell}(1 + v\mathbb{A}_0), \tag{2.4}$$

$$\operatorname{ord}(F_{b,b'}) > 1 - \ell' \text{ if } \ell < \ell' \text{ and } b' \neq \widetilde{f}b, \tag{2.5}$$

$$\operatorname{ord}(E_{b,b'}) > -\ell' \text{ if } \ell < \ell' + 1 \text{ and } b' \neq \widetilde{e}b. \tag{2.6}$$

Theorem 2.14 ([EK2, Theorem 4.1, Corollary 4.4])**.** *Assume the conditions (2.1)–(2.6). Let L be the $\mathbf{A}_0$-submodule $\sum_{b \in B} \mathbf{A}_0 G(b)$ of M. Then we have $\widetilde{e}L \subset L$ and $\widetilde{f}L \subset L$. Moreover we have*

$$\widetilde{e}G(b) \equiv G(\widetilde{e}b) \mod vL, \quad \widetilde{f}G(b) \equiv G(\widetilde{f}b) \mod vL$$

for any $b \in B$. Here we understand $G(0) = 0$.

In [EK2], this theorem is proved under more general assumptions.

2.4. **Perverse Sheaves.**

2.4.1. *Perverse Sheaves.* In this paper, we consider algebraic varieties over $\mathbb{C}$. Let $\mathscr{D}(X)$ be the bounded derived category of constructible complexes of sheaves on an algebraic variety X. We denote by $\mathscr{D}^{\leq 0}(X)$ (resp. $\mathscr{D}^{\geq 0}(X)$) the full subcategory of $\mathscr{D}(X)$ consisting of objects L satisfying $H^k(L) = 0$ for $k > 0$ (resp. $k < 0$). Put $\mathscr{D}^{\leq n} = \mathscr{D}^{\leq 0}[-n]$ and $\mathscr{D}^{\geq n} = \mathscr{D}^{\geq 0}[-n]$.
For a morphism $f\colon X \to Y$ of algebraic varieties X and Y, let f^* be the inverse image, $f_!$ the direct image with proper support and $D\colon \mathscr{D}(X) \to \mathscr{D}(X)$ the Verdier duality functor.

Lemma 2.15.

(i) *Suppose that $f\colon X \to Y$ is smooth with the fiber dimension d. Then $D(f^*L) \cong f^*D(L)[2d]$ for $L \in \mathscr{D}(Y)$.*
(ii) *Suppose that $f\colon X \to Y$ is proper. Then $D(f_!L) \cong f_!D(L)$ for $L \in \mathscr{D}(X)$.*

Let $({}^p\mathscr{D}^{\leq 0}(X), {}^p\mathscr{D}^{\geq 0}(X))$ be the perverse t-structure and $\operatorname{Perv}(X) := {}^p\mathscr{D}^{\leq 0}(X) \cap {}^p\mathscr{D}^{\geq 0}(X)$.

Lemma 2.16. *Suppose $L \in {}^p\mathscr{D}^{\leq 0}(X)$ and $K \in {}^p\mathscr{D}^{\geq 0}(X)$, then $H^j(\mathbf{R}\mathscr{H}om(L, K)) = 0$ for $j < 0$, namely $\mathbf{R}\mathscr{H}om(L, K) \in \mathscr{D}^{\geq 0}(X)$.*

Let ${}^pH^k(\ )$ be the k-th perverse cohomology sheaf. We say that an object L in $\mathscr{D}(X)$ is semisimple if L is isomorphic to the direct sum $\oplus_k {}^pH^k(L)[-k]$ and if each ${}^pH^k(L)$ is a semisimple perverse sheaf. Assume that we are given an action of a connected algebraic group G on X. A semisimple object L in $\mathscr{D}(X)$ is said to be G-equivariant if each ${}^pH^i(L)$ is a G-equivariant perverse sheaf.

Lemma 2.17.

(i) *Suppose that $f\colon X \to Y$ is smooth with connected fibers of dimension d. Then we have a fully faithful functor $\operatorname{Perv}(Y) \to \operatorname{Perv}(X)$ given by $K \mapsto f^*K[d]$. Moreover if K is simple, then $f^*K[d]$ is simple.*

(ii) Let G be a connected algebraic group of dimension d and $\mathrm{Perv}_G(X)$ the category of G-equivariant perverse sheaves. Suppose that $f\colon X \to Y$ is a principal G-bundle. The functors

$$\mathrm{Perv}(Y) \to \mathrm{Perv}_G(X)\colon K \mapsto f^*K[d]$$

and

$$\mathrm{Perv}_G(X) \to \mathrm{Perv}(Y)\colon L \mapsto ({}^pH^{-d}f_*L)$$

*define an equivalence of categories, quasi-inverse to each other.
Moreover if K is a semisimple object of $\mathscr{D}(Y)$, then f^*K is a G-equivariant semisimple object in $\mathscr{D}(X)$. Conversely, if L is a G-equivariant semisimple object of $\mathscr{D}(X)$, then there is a unique semisimple object $K \in \mathscr{D}(Y)$ such that $L \cong f^*K$.*

We denote by $\mathbf{1}_X$ the constant sheaf on X.

Lemma 2.18 ([BBD], [Lus3])**.**

(1) *Let $f\colon X \to Y$ be a projective morphism with X smooth. Then $f_!\mathbf{1}_X \in \mathscr{D}(Y)$ is semisimple.*

(2) *Let $f\colon X \to Y$ be a morphism. Assume that there exists a partition $X = X_0 \sqcup X_1 \sqcup \cdots \sqcup X_m$ such that $X_{\leq j} = X_0 \cup X_1 \cup \cdots \cup X_j$ is closed for $j = 0, 1, \ldots, m$. Assume that, for each j, the restriction $f_j\colon X_j \to Y$ of f decomposes as $X_j \xrightarrow{f''_j} Z_j \xrightarrow{f'_j} Y$ such that Z_j is smooth, f''_j is an affine bundle and f'_j is projective. Then $f_!\mathbf{1}_X \in \mathscr{D}(Y)$ is semisimple. Moreover, we have $f_!\mathbf{1}_X \cong \bigoplus_j (f_j)_!\mathbf{1}_{X_j}$.*

2.4.2. *Simple objects.* Let Y be an irreducible variety and U a Zariski open subset of Y. Set $Z := Y \backslash U$ and $i\colon Z \hookrightarrow Y$.

Proposition 2.19. *For $F \in \mathrm{Perv(U)}$, there exists a unique perverse sheaf ${}^\pi F$ on Y satisfying*

(i) ${}^\pi F|_U \cong F$,
(ii) $i^*({}^\pi F) \in {}^p\mathscr{D}^{\leq -1}(Z)$,
(iii) $i^!({}^\pi F) \in {}^p\mathscr{D}^{\geq 1}(Z)$.

We call ${}^\pi F$ the minimal extension of F. We have the following properties of the minimal extension:

(1) ${}^\pi F$ has neither non-trivial subobject nor non-trivial quotient object whose support is contained in Z.
(2) If F is simple, then ${}^\pi F$ is simple.
(3) For the Verdier duality functors D_Y and D_U, we have $D_Y({}^\pi F) \cong {}^\pi(D_U(F))$.

Let X be a variety, Y an irreducible locally closed smooth subvariety of X. For a simple local system L on Y, the minimal extension ${}^\pi L[\dim Y]$ is called the intersection cohomology complex of Y. We can regard ${}^\pi L[\dim Y]$ as a simple perverse on X whose support is the closure $\overline{Y}$ of Y. Conversely, any simple object in Perv(X) is obtained in this way.

Theorem 2.20 ([BBD])**.** *For a simple perverse sheaf F on X, there exist an irreducible closed subvariety Y and an simple local system L on Y such that $F \cong {}^\pi L[\dim Y]$. Moreover, for simple perverse sheaves F_1 and F_2, we have $\mathrm{Ext}^0(\mathrm{F}_1, \mathrm{F}_2) = \mathrm{Hom}_{\mathrm{Perv(X)}}(\mathrm{F}_1, \mathrm{F}_2) = \mathbb{C}$ or 0 according that F_1 and F_2 are isomorphic or not.*

2.4.3. *Fourier-Sato-Deligne transforms.* Let $E \to S$ be a vector bundle and $E^* \to S$ the dual vector bundle. Hence $\mathbb{C}^\times$ acts on E and E^*. We say that $L \in \mathscr{D}(E)$ is monodromic if $H^j(L)$ is locally constant on every $\mathbb{C}^*$-orbit of E. Let $\mathscr{D}_{\mathrm{mono}}(E)$ be the full subcategory of $\mathscr{D}(E)$ consisting of monodromic objects. Then we can define the Fourier transform

$$\Phi_{E/S} \colon \mathscr{D}_{\mathrm{mono}}(E) \to \mathscr{D}_{\mathrm{mono}}(E^*).$$

We will use the following properties of Φ.

Proposition 2.21 (e.g. [KS], [Lau])**.**

1. *For $K \in \mathscr{D}_{\mathrm{mono}}(E)$, we have $\Phi_{E^*/S} \circ \Phi_{E/S}(K) \cong a^*K$, where $a \colon E \to E$ is the multiplication by -1 on each fiber of E.*
2. *For a perverse sheaf $K \in \mathscr{D}_{\mathrm{mono}}(E)$, $\Phi_{E/S}(K)$ is a perverse sheaf in $\mathscr{D}_{\mathrm{mono}}(E^*)$.*
3. *Let E_1 and E_2 be two vector bundles over S with rank r_1 and r_2. Let $f \colon E_1 \to E_2$ be a morphism of vector bundles and ${}^tf \colon E_2^* \to E_1^*$ the transpose of f. Then we have*
$$\Phi_{E_2/S} \circ f_! \cong ({}^tf)^* \circ \Phi_{E_1/S}[r_2 - r_1], \quad ({}^tf)_! \circ \Phi_{E_2/S} \cong \Phi_{E_1/S} \circ f^*[r_1 - r_2].$$
4. *Suppose that $E_1 \to S_1$ and $E \to S$ are two vector bundles. If the following two diagrams*
$$\begin{tikzcd} E_1 \arrow[r, "f_E"] \arrow[d] & E \arrow[d] \\ S_1 \arrow[r, "\rho"] & S \end{tikzcd} \qquad \begin{tikzcd} E_1^* \arrow[r, "f_{E^*}"] \arrow[d] & E^* \arrow[d] \\ S_1 \arrow[r, "\rho"] & S \end{tikzcd}$$
are Cartesian, then we have
$$\Phi_{E/S} \circ (f_E)_! \cong (f_{E^*})_! \circ \Phi_{E_1/S_1}, \quad \Phi_{E_1/S_1} \circ (f_E)^* \cong (f_{E^*})^* \circ \Phi_{E/S}.$$
5. *The Fourier transforms commute with the Verdier duality functors.*

2.5. **Quivers.**

Let I and α_i's be as in 2.1.

Definition 2.22. *A quiver (I, H) associated with the symmetric Cartan matrix is a following data:*

- (i) *a set H,*
- (ii) *two maps* out, in: $H \to I$ *such that* $\mathrm{out}(h) \neq \mathrm{in}(h)$ *for any $h \in H$,*
- (iii) *an involution $h \mapsto \bar{h}$ on H satisfying* $\mathrm{out}(\bar{h}) = \mathrm{in}(h)$ *and* $\mathrm{in}(\bar{h}) = \mathrm{out}\, h$,
- (iv) $\sharp\{h \in H \,|\, \mathrm{out}(h) = i, \mathrm{in}(h) = j\} = -(\alpha_i, \alpha_j)$ *for $i \neq j$.*

An orientation of a quiver (I, H) is a subset Ω of H such that $\Omega \cap \overline{\Omega} = \phi$ and $\Omega \cup \overline{\Omega} = H$. For a fixed orientation Ω, we call a vertex $i \in I$ a sink if $\mathrm{out}(h) \neq i$ *for any $h \in \Omega$.*

Definition 2.23. *Let $\mathcal{V}$ be the category of I-graded vector spaces $\mathbf{V} = (\mathbf{V}_i)_i$ with morphisms being linear maps respecting the grading. Put* $\mathrm{wt}(\mathbf{V}) = \sum_{i \in I} (\dim \mathbf{V}_i)\alpha_i$.

Let $\mathbf{S}_i$ be an I-graded vector space such that $\mathrm{wt}(\mathbf{S}_i) = \alpha_i$.

Definition 2.24. *For $\mathbf{V} \in \mathcal{V}$ and a subset Ω of H, we define*

$$\mathbf{E}_{\mathbf{V},\Omega} \colon = \bigoplus_{h \in \Omega} \mathrm{Hom}(\mathbf{V}_{\mathrm{out}(h)}, \mathbf{V}_{\mathrm{in}(h)}).$$

The algebraic group $\mathbf{G}_{\mathbf{V}} = \prod_{i \in I} GL(\mathbf{V}_i)$ acts on $\mathbf{E}_{\mathbf{V},\Omega}$ by $(g, x) \mapsto gx$ where $(gx)_h = g_{\mathrm{in}(h)} x_h g_{\mathrm{out}(h)}^{-1}$.
The group $(\mathbb{C}^\times)^\Omega$ also acts on $\mathbf{E}_{\mathbf{V},\Omega}$ by $x_h \mapsto c_h x_h$ $(h \in \Omega, c_h \in \mathbb{C}^\times)$.
For $x \in \mathbf{E}_{\mathbf{V},\Omega}$, an I-graded subspace $\mathbf{W} \subset \mathbf{V}$ is x-stable if $x_h(\mathbf{W}_{\mathrm{out}(h)}) \subset \mathbf{W}_{\mathrm{in}(h)}$ for any $h \in \Omega$.

Note that $E_{\mathbf{S}_i,\Omega} \cong \{\mathrm{pt}\}$.

3. A Review on Lusztig's Geometric Construction

We give a quick review on Lusztig's theory in [Lus1] and [Lus2] (cf. [Lus3]). For a sequence $\mathbf{i} = (i_1, \dots, i_m) \in I^m$ and a sequence $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{Z}^m_{\geq 0}$, a flag of type $(\mathbf{i}, \mathbf{a})$ is by definition a finite decreasing sequence $F = (\mathbf{V} = \mathbf{F}^0 \supset \mathbf{F}^1 \supset \cdots \supset \mathbf{F}^m = \{0\})$ of I-graded subspaces of $\mathbf{V}$ such that the I-graded vector space $\mathbf{F}^{\ell-1}/\mathbf{F}^{\ell}$ vanishes in degrees $\neq i_\ell$ and has dimension a_ℓ in degree i_ℓ. We denote by $\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}$ the set of pairs (x, F) such that $x \in \mathbf{E}_{\mathbf{V},\Omega}$ and F is an x-stable flag of type $(\mathbf{i}, \mathbf{a})$. The group $G_{\mathbf{V}}$ acts on $\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}$. The first projection $\pi_{\mathbf{i},\mathbf{a}}\colon \widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega} \to \mathbf{E}_{\mathbf{V},\Omega}$ is a $G_{\mathbf{V}}$-equivariant projective morphism.
By Lemma 2.18, $L_{\mathbf{i},\mathbf{a};\Omega}\colon = (\pi_{\mathbf{i},\mathbf{a}})_!(\mathbf{1}_{\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}}) \in \mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$ is a semisimple complex. We define $\mathscr{P}_{\mathbf{V},\Omega}$ as the set of the isomorphism classes of simple perverse sheaves $L \in \mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$ satisfying the following property: L appears as a direct summand of $L_{\mathbf{i},\mathbf{a};\Omega}[d]$ for some d and $(\mathbf{i}, \mathbf{a})$. We denote by $\mathscr{Q}_{\mathbf{V},\Omega}$ the full subcategory of $\mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$ consisting of all objects which are isomorphic to finite direct sums of complexes of the form $L[d]$ for various $L \in \mathscr{P}_{\mathbf{V},\Omega}$ and various integers d. Any complex in $\mathscr{P}_{\mathbf{V},\Omega}$ is $G_{\mathbf{V}} \times (\mathbb{C}^\times)^\Omega$-equivariant.
Let $\mathbf{T}, \mathbf{W}, \mathbf{V}$ be I-graded vector spaces such that $\mathrm{wt}(\mathbf{V}) = \mathrm{wt}(\mathbf{W}) + \mathrm{wt}(\mathbf{T})$. We consider the following diagram

$$\begin{tikzcd} \mathbf{E}_{\mathbf{T},\Omega} \times \mathbf{E}_{\mathbf{W},\Omega} & \mathbf{E}'_\Omega \arrow[l, "p_1"'] \arrow[r, "p_2"] & \mathbf{E}''_\Omega \arrow[r, "p_3"] & \mathbf{E}_{\mathbf{V},\Omega}. \end{tikzcd}$$

Here $\mathbf{E}''_\Omega$ is the variety of (x, W) where $x \in \mathbf{E}_{\mathbf{V},\mathbf{\Omega}}$ and W is an x-stable I-graded subspace of $\mathbf{V}$ such that $\mathrm{wt}\, W = \mathrm{wt}\, \mathbf{W}$. The variety $\mathbf{E}'_\Omega$ consists of $(x, W, \varphi^{\mathbf{W}}, \varphi^{\mathbf{T}})$ where $(x, W) \in \mathbf{E}''_\Omega$, $\varphi^{\mathbf{W}}\colon \mathbf{W} \cong W$, and $\varphi^{\mathbf{T}}\colon \mathbf{T} \cong \mathbf{V}/W$. The morphisms p_1, p_2 and p_3 are given by $p_1(x, W, \varphi^{\mathbf{W}}, \varphi^{\mathbf{T}}) = (x|_{\mathbf{T}}, x|_{\mathbf{W}})$, $p_2(x, W, \varphi^{\mathbf{W}}, \varphi^{\mathbf{T}}) = (x, W)$ and $p_3(x, W) = x$. Then p_1 is smooth with connected fibers, p_2 is a principal $G_{\mathbf{T}} \times G_{\mathbf{W}}$-bundle, and p_3 is projective. For a $G_{\mathbf{T}}$-equivariant semisimple complex $K_{\mathbf{T}}$ and a $G_{\mathbf{W}}$-equivariant semisimple complex $K_{\mathbf{W}}$, there exists a unique semisimple complex K'' satisfying $p_1^*(K_{\mathbf{T}} \boxtimes K_{\mathbf{W}}) = p_2^* K''$. We define $K_{\mathbf{T}} * K_{\mathbf{W}}\colon = (p_3)_!(K'') \in \mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$.
For an I-graded subspace $\mathbf{U}$ of $\mathbf{V}$ such that $\mathbf{V}/\mathbf{U} \cong \mathbf{T}$, we also consider the following diagram

$$\begin{tikzcd} \mathbf{E}_{\mathbf{T},\Omega} \times \mathbf{E}_{\mathbf{U},\Omega} & \mathbf{E}(\mathbf{U}, \mathbf{V})_\Omega \arrow[l, "p"'] \arrow[r, hook, "\iota"] & \mathbf{E}_{\mathbf{V},\Omega}. \end{tikzcd}$$

Here $\mathbf{E}(\mathbf{U}, \mathbf{V})_\Omega$ is the variety of $x \in \mathbf{E}_{\mathbf{V},\Omega}$ such that $\mathbf{U}$ is x-stable. For $K \in \mathscr{D}(\mathbf{E}_{\mathbf{V},\Omega})$, we define $\mathrm{Res}_{\mathbf{T},\mathbf{U}}(K)\colon = p_! \iota^*(K)$.
We define $K_{\mathbf{V},\Omega}$ as the Grothendieck group of $\mathscr{Q}_{\mathbf{V},\Omega}$. It is the additive group generated by the isomorphism classes (L) of objects $L \in \mathscr{Q}_{\mathbf{V},\Omega}$ with the relation $(L) = (L') + (L'')$ when $L \cong L' \oplus L''$. The group $K_{\mathbf{V},\Omega}$ has a $\mathbb{Z}[v, v^{-1}]$-module structure by $v(L) = (L[1])$ and $v^{-1}(L) = (L[-1])$ for $L \in \mathscr{Q}_{\mathbf{V},\Omega}$. Hence, $K_{\mathbf{V},\Omega}$ is a free $\mathbb{Z}[v, v^{-1}]$-module with a basis $\{(L) | L \in \mathscr{P}_{\mathbf{V},\Omega}\}$. We define $K_\Omega\colon = \bigoplus_{\mathbf{V}} K_{\mathbf{V},\Omega}$ where $\mathbf{V}$ runs over the isomorphism classes of I-graded vector spaces. Recall that $\mathbf{S}_i$ is an I-graded vector space such that $\mathrm{wt}(\mathbf{S}_i) = \alpha_i$. Then we can define the induction $f_i\colon K_{\mathbf{W},\Omega} \to K_{\mathbf{V},\Omega}$ and the restriction $e'_i\colon K_{\mathbf{V},\Omega} \to K_{\mathbf{W},\Omega}$ by

$$f_i(K)\colon = v^{\dim \mathbf{W}_i + \sum_{i \xrightarrow{\Omega} j} \dim \mathbf{W}_j} (\mathbf{1}_{\mathbf{S}_i} * K), \quad e'_i(K)\colon = v^{-\dim \mathbf{W}_i + \sum_{i \xrightarrow{\Omega} j} \dim \mathbf{W}_j} \mathrm{Res}_{\mathbf{S}_i,\mathbf{V}}(K).$$

Then Lusztig's main theorem is stated as follows.

Theorem 3.1 (Lusztig).

(i) The operators e_i' and f_i define the action of the reduced v-analogue $B_v(\mathfrak{g})$ of $\mathfrak{g}$ on $K_\Omega \otimes_{\mathbb{Z}[v,v^{-1}]} \mathbb{Q}(v)$. The $B_v(\mathfrak{G})$-module $K_\Omega \otimes_{\mathbb{Z}[v,v^{-1}]} \mathbb{Q}(v)$ is isomorphic to $U_v^-(\mathfrak{g})$. The involution induced by the Verdier duality functor coincides with the bar involution on $U_v^-(\mathfrak{g})$.

(ii) The simple perverse sheaves in $\sqcup_{\mathbf{V}} \mathscr{P}_{\mathbf{V},\Omega}$ give a lower global basis of $U_v^-(\mathfrak{g})$.

4. Quivers with an Involution θ

4.1. **Quivers with an involution θ.**

Definition 4.1. *A θ-quiver is a data:*

(1) a quiver (I,H),

(2) involutions $\theta\colon I \to I$ and $\theta\colon H \to H$,

satisfying

(a) $\operatorname{out}(\theta(h)) = \theta(\operatorname{in}(h))$ *and* $\operatorname{in}(\theta(h)) = \theta(\operatorname{out}(h))$,

(b) If $\theta(\operatorname{out}(h)) = \operatorname{in}(h)$, *then* $\theta(h) = h$,

(c) $\theta(\overline{h}) = \overline{\theta(h)}$,

(d) There is no $i \in I$ such that $\theta(i) = i$

A θ-orientation is an orientation of (I,H) such that Ω is stable by θ.

From the assumption (d), any vertex i is a sink with respect to some θ-orientation Ω.

Example 4.2. We give two θ-orientations for the case of Example 2.13. The vertex 1 is a sink in the right example.



,



.

Example 4.3. Our definition of a θ-quiver contains the case of type $A_1^{(1)}$. The following three figures are three θ-orientations in this case.



Definition 4.4. *A θ-symmetric I-graded vector space $\mathbf{V}$ is an I-graded vector space endowed with a non-degenerate symmetric bilinear form $(\,\bullet\,,\bullet\,)\colon \mathbf{V} \times \mathbf{V} \to \mathbb{C}$ such that $\mathbf{V}_i$ and $\mathbf{V}_j$ are orthogonal if $j \neq \theta(i)$. For an I-graded subspace $\mathbf{W}$ of $\mathbf{V}$, we set*

$$\mathbf{W}^\perp\colon = \{v \in \mathbf{V} \mid (v,w) = 0 \text{ for any } w \in \mathbf{W}\}.$$

Hence $(\mathbf{W}^\perp)_{\theta(i)} \cong (\mathbf{V}_i/\mathbf{W}_i)^*$.

Note that if $\mathbf{W} \supset \mathbf{W}^\perp$, then $\mathbf{W}/\mathbf{W}^\perp$ has a structure of θ-symmetric I-graded vector space. Note that two θ-symmetric I-graded vector spaces with the same dimension are isomorphic.

Definition 4.5. *Let (I,H) be a θ-quiver. For a θ-symmetric I-graded vector space $\mathbf{V}$ and a θ-stable subset Ω of H, we define*

$${}^\theta\mathbf{E}_{\mathbf{V},\Omega}\colon = \{x \in \mathbf{E}_{\mathbf{V},\Omega} \mid x_{\theta(h)} = -{}^t x_h \in \operatorname{Hom}(\mathbf{V}_{\theta(\operatorname{in}(h))}, \mathbf{V}_{\theta(\operatorname{out}(h))}) \text{ for any } h \in \Omega\}.$$

The algebraic group ${}^\theta\mathbf{G}_{\mathbf{V}}\colon = \{g \in \mathbf{G}_{\mathbf{V}} \mid {}^t g_i^{-1} = g_{\theta(i)} \text{ for any } i\}$ naturally acts on ${}^\theta\mathbf{E}_{\mathbf{V},\Omega}$. Set $(\mathbb{C}^\times)^{\Omega,\theta}\colon = \{(c_h)_{h\in\Omega} \mid c_h \in \mathbb{C}^\times \text{ and } c_{\theta(h)} = c_h\}$. The group $(\mathbb{C}^\times)^{\Omega,\theta}$ also acts on ${}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ by $x_h \mapsto c_h x_h$ $(h \in \Omega)$. These two actions commute with each other.

Definition 4.6. *For a θ-symmetric I-graded vector space $\mathbf{V}$, a sequence $\mathbf{i} = (i_1, \ldots, i_{2m}) \in I^{2m}$ such that $\theta(i_\ell) = i_{2m-\ell+1}$ and a sequence $\mathbf{a} = (a_1, \ldots, a_{2m}) \in \mathbb{Z}^m_{\geq 0}$ such that $a_{2m-\ell+1} = a_\ell$, we say that a flag of I-graded subspace of $\mathbf{V}$*

$$F = (\mathbf{V} = \mathbf{F}^0 \supset \mathbf{F}^1 \supset \cdots \supset \mathbf{F}^m \supset \mathbf{F}^{m+1} \supset \cdots \supset \mathbf{F}^{2m} = \{0\})$$

is of type $(\mathbf{i}, \mathbf{a})$ if

(i) $\dim(\mathbf{F}^{\ell-1}/\mathbf{F}^\ell)_i = \begin{cases} a_\ell & (i = i_\ell) \\ 0 & (i \neq i_\ell) \end{cases}$,

(ii) $\mathbf{F}^{2m-\ell} = (\mathbf{F}^\ell)^\perp$.

Then we have $\mathrm{wt}\, \mathbf{V} = \sum_{1 \leq \ell \leq 2m} a_\ell \alpha_{i_\ell}$. *We denote by ${}^\theta\mathcal{F}_{\mathbf{i},\mathbf{a}}$ the set of flags of type $(\mathbf{i}, \mathbf{a})$. For $x \in {}^\theta\mathbf{E}_{\mathbf{V},\Omega}$, a flag F of type $(\mathbf{i}, \mathbf{a})$ is x-stable if $\mathbf{F}^\ell$ $(\ell = 1, \ldots, 2m)$ are x-stable. We define*

$${}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega} \colon = \{(x, F) \in {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \times {}^\theta\mathcal{F}_{\mathbf{i},\mathbf{a}} \mid F \text{ is } x\text{-stable}\}.$$

The group ${}^\theta\mathbf{G}_\mathbf{V}$ naturally acts on ${}^\theta\mathcal{F}_{\mathbf{i},\mathbf{a}}$ and ${}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}$.

Note that $x \colon \mathbf{V} \to \mathbf{V} \cong \mathbf{V}^*$ in ${}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ may be regarded as a skew-symmetric form on $\mathbf{V}$, and the condition that F is x-stable is equivalent to the one $x(\mathbf{F}^\ell, \mathbf{F}^{2m-\ell}) = 0$ for any ℓ. The following lemma is obvious.

Lemma 4.7. *The variety ${}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}$ is smooth and irreducible. The first projection ${}^\theta\pi_{\mathbf{i},\mathbf{a}} \colon {}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega} \to {}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ is ${}^\theta\mathbf{G}_\mathbf{V} \times (\mathbb{C}^\times)^{\Omega,\theta}$-equivariant and projective.*

4.2. **Perverse sheaves on ${}^\theta\mathbf{E}_{\mathbf{V},\Omega}$.** Let Ω be a θ-orientation. By Lemma 4.7 and Lemma 2.18,

$${}^\theta L_{\mathbf{i},\mathbf{a};\Omega} \colon = ({}^\theta\pi_{\mathbf{i},\mathbf{a}})_!(\mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}})$$

is a semisimple complex in $\mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$.

Definition 4.8. *We define ${}^\theta\mathscr{P}_{\mathbf{V},\Omega}$ as the set of the isomorphism classes of simple perverse sheaves L in $\mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$ satisfying the property: L appears in ${}^\theta L_{\mathbf{i},\mathbf{a};\Omega}[d]$ as a direct summand for some integer d and $(\mathbf{i}, \mathbf{a})$. We denote by ${}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$ the full subcategory of $\mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$ consisting of objects which are isomorphic to finite direct sums of $L[d]$ with $L \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$ and $d \in \mathbb{Z}$.*

Note that any object in ${}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$ is ${}^\theta\mathbf{G}_\mathbf{V} \times (\mathbb{C}^\times)^{\Omega,\theta}$-equivariant.

4.3. **Multiplications and Restrictions.** Fix θ-symmetric and I-graded vector spaces $\mathbf{V}$ and $\mathbf{W}$, and an I-graded vector space $\mathbf{T}$ such that $\mathrm{wt}(\mathbf{V}) = \mathrm{wt}(\mathbf{W}) + \mathrm{wt}(\mathbf{T}) + \theta(\mathrm{wt}(\mathbf{T}))$. We consider the following diagram

$$\begin{tikzcd} \mathbf{E}_{\mathbf{T},\Omega} \times {}^\theta\mathbf{E}_{\mathbf{W},\Omega} & {}^\theta\mathbf{E}'_\Omega \arrow[l, "p_1"'] \arrow[r, "p_2"] & {}^\theta\mathbf{E}''_\Omega \arrow[r, "p_3"] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \end{tikzcd} .$$

Here ${}^\theta\mathbf{E}''_\Omega$ is the variety of (x, V) where $x \in {}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ and V is an x-stable I-graded subspace of $\mathbf{V}$ such that $V \supset V^\perp$ and $\mathrm{wt}(\mathbf{V}/V) = \mathrm{wt}(\mathbf{T})$, and we denote by ${}^\theta\mathbf{E}'_\Omega$ the variety of $(x, V, \varphi^\mathbf{W}, \varphi^\mathbf{T})$ where $(x, V) \in {}^\theta\mathbf{E}''_\Omega$, $\varphi^\mathbf{W} \colon \mathbf{W} \xrightarrow{\sim} V/V^\perp$ is an isomorphism of θ-symmetric I-graded vector spaces and $\varphi^\mathbf{T} \colon \mathbf{T} \xrightarrow{\sim} \mathbf{V}/V$ is an isormorphism of I-graded vector spaces. We define p_1, p_2 and p_3 by $p_1(x, V, \varphi^\mathbf{W}, \varphi^\mathbf{T}) = (x^\mathbf{T}, x^\mathbf{W})$, $p_2(x, V, \varphi^\mathbf{W}, \varphi^\mathbf{T}) = (x, V)$ and $p_3(x, V) = x$. Here the morphism $x^\mathbf{W}, x^\mathbf{T}$ are defined by

$$x^\mathbf{W}_h = {\varphi^\mathbf{W}_{\mathrm{in}(h)}}^{-1} \circ (x|_{V/V^\perp})_h \circ \varphi^\mathbf{W}_{\mathrm{out}(h)}, \quad x^\mathbf{T}_h = {\varphi^\mathbf{T}_{\mathrm{in}(h)}}^{-1} \circ (x|_{\mathbf{V}/V})_h \circ \varphi^\mathbf{T}_{\mathrm{out}(h)}.$$

Then p_1 is smooth with connected fibers, p_2 is a principal $\mathbf{G_T} \times {}^\theta\mathbf{G_W}$-bundle and p_3 is projective.
For a $\mathbf{G_T}$-equivariant semisimple object $K_\mathbf{T} \in \mathscr{Q}_{\mathbf{T},\Omega}$ and a ${}^\theta\mathbf{G_W}$-equivariant semisimple object $K_\mathbf{W} \in {}^\theta\mathscr{Q}_{\mathbf{W},\Omega}$, there exists a unique semisimple object $K'' \in \mathscr{D}({}^\theta\mathbf{E}''_\Omega)$ satisfying $p_1^*(K_\mathbf{T} \boxtimes K_\mathbf{W}) = p_2^* K''$.

Definition 4.9. *We define* $K_\mathbf{T} * K_\mathbf{W} \colon = (p_3)_!(K'') \in \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$.

Next, we fix an I-graded vector space U such that

$$\mathbf{V} \supset U \supset U^\perp \supset \{0\}.$$

We also fix an isomorphism $\mathbf{W} \cong U/U^\perp$ as θ-symmetric I-graded vector spaces and an isomorphism $\mathbf{T} \cong \mathbf{V}/U$ as I-graded vector spaces. We consider the following diagram

$$\begin{tikzcd}
\mathbf{E}_{\mathbf{T},\Omega} \times {}^\theta\mathbf{E}_{\mathbf{W},\Omega} & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega \arrow[l, "p"'] \arrow[r, hook, "\iota"] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega}
\end{tikzcd}$$

where

$${}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega = \{x \in {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \mid U \text{ is } x\text{-stable}\}$$

and $p(x) = (x^\mathbf{T}, x^\mathbf{W})$, $\iota(x) = x$.

Definition 4.10. *For* $K \in \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$, *we define* $\mathrm{Res}_{\mathbf{T},\mathbf{W}}(K) \colon = p_! \iota^*(K)$.

Proposition 4.11. *Let* $\mathbf{V}$ *and* $\mathbf{W}$ *be* θ*-symmetric* I*-graded vector spaces such that* $\mathrm{wt}\,\mathbf{V} = \mathrm{wt}\,\mathbf{W} + \alpha_i + \alpha_{\theta(i)}$. *For* $a \in \mathbb{Z}_{\geq 0}$, *let* $\mathbf{S}_i^a$ *be an* I*-graded vector space such that* $\mathrm{wt}(\mathbf{S}_i^a) = a\alpha_i$.

(i) *Suppose* ${}^\theta L_{\mathbf{i},\mathbf{a};\Omega} \in \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{W},\Omega})$. *We have*

$$\mathbf{1}_{\mathbf{S}_i^a} * {}^\theta L_{\mathbf{i},\mathbf{a};\Omega} = L_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}.$$

for $a \in \mathbb{Z}_{\geq 0}$.

(ii) *Suppose* ${}^\theta L_{\mathbf{i},\mathbf{a};\Omega} \in \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$ *and* $a_\ell > 0$ *for all* ℓ *such that* $i_\ell = i$. *For* $1 \leq k \leq 2m$ *such that* $i_k = i$, *we define* $\mathbf{a}^{(k)} = (a_1^{(k)}, \cdots, a_{2m}^{(k)})$ *by* $a_\ell^{(k)} = a_\ell - \delta_{\ell,k} - \delta_{\ell,2m-k+1}$ *and we set*

$$M_k(\mathbf{i},\mathbf{a}^{(k)}) = \sum_{i_\ell = i, \ell < k} a_\ell^{(k)} + \sum_{k < \ell, h \in \Omega; \mathrm{out}(h) = i, \mathrm{in}(h) = i_\ell} a_\ell^{(k)}.$$

Then we have

$$\mathrm{Res}_{\mathbf{S}_i,\mathbf{W}}({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) = \bigoplus_{i_k = i} {}^\theta L_{\mathbf{i},\mathbf{a}^{(k)};\Omega}[-2M_k(\mathbf{i},\mathbf{a}^{(k)})].$$

Proof. (1) We consider the following diagram:

$$\begin{tikzcd}
{}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega} \arrow[d, "{}^\theta\pi_{\mathbf{i},\mathbf{a}}"'] \arrow[dr, phantom, "\square"] & {}^\theta\widetilde{E} \arrow[l, "p_1'"'] \arrow[r, "p_2'"] \arrow[d, "\rho'"'] \arrow[dr, phantom, "\square"] & {}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a);\Omega} \arrow[d, "\rho''"'] \arrow[dr, "\pi_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}"] & \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega} & {}^\theta\mathbf{E}'_\Omega \arrow[l, "p_1"'] \arrow[r, "p_2"] & {}^\theta\mathbf{E}''_\Omega \arrow[r, "p_3"] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega}
\end{tikzcd}$$

where

$${}^\theta\widetilde{E} \colon = \{(x, F, \varphi^\mathbf{W}) \mid (x, F) \in {}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}, \varphi^\mathbf{W} \colon \mathbf{W} \cong \mathbf{F}^1/\mathbf{F}^{2m+1}\}.$$

Here, $\rho'' \colon {}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)} \to {}^\theta\mathbf{E}''_\Omega$ is given by $(x, F) \mapsto (x, \mathbf{F}^1)$. Then ρ'' is projective and $p_3 \circ \rho'' = {}^\theta\pi_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}$. Hence $\rho''_!(\mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}})$ is semisimple and ${}^\theta L_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a);\Omega} = (\pi_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)})_!(\mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}}) = (p_3)_!(\rho'')_!(\mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}})$. On the other hand, we have

$$p_2^*(\rho''_! \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}}) = \rho'_!(p'_2)^* \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}} = \rho'_!(p'_1)^* \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}} = p_1^*({}^\theta\pi_{\mathbf{i},\mathbf{a}})_! \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}} = p_1^*({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}).$$

Hence we have $\mathbf{1}_{\mathbf{S}_i^a} * {}^\theta L_{\mathbf{i},\mathbf{a};\Omega} = (p_3)_! \rho''_!(\mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}}) = {}^\theta L_{(i,\mathbf{i},\theta(i)),(a,\mathbf{a},a)}$.

(2) Set ${}^\theta\widetilde{\mathcal{F}}(\mathbf{W},\mathbf{V}) = \{(x,F) \in {}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega} \mid U \text{ is } x\text{-stable}\}$ and ${}^\theta\mathcal{F}^{(k)}_{\mathbf{i},\mathbf{a};\Omega} = \{F \in {}^\theta\mathcal{F}_{\mathbf{i},\mathbf{a};\Omega} \mid \mathbf{F}^k \subset U, \mathbf{F}^{k-1} \not\subset U\}$. We define

$$ {}^\theta\widetilde{\mathcal{F}}_k(\mathbf{W},\mathbf{V}) \colon = \{(x,F) \in {}^\theta\widetilde{\mathcal{F}}(\mathbf{W},\mathbf{V}) \mid F \in {}^\theta\mathcal{F}^{(k)}_{\mathbf{i},\mathbf{a};\Omega}\}. $$

Then the locally closed smooth subvarieties ${}^\theta\widetilde{\mathcal{F}}_k(\mathbf{W},\mathbf{V})$ $(1 \le k \le 2m, i_k = i)$ give a partition ${}^\theta\widetilde{\mathcal{F}}(\mathbf{W},\mathbf{V})$.

For a flag F of $\mathbf{V}$, we define the flag $F|_{U/U^\perp}$ by

$$ F|_{U/U^\perp} = (U/U^\perp = (\mathbf{F}^0 \cap U)/(\mathbf{F}^0 \cap U^\perp) \supset \cdots \supset (\mathbf{F}^{2m} \cap U)/(\mathbf{F}^{2m} \cap U^\perp) = \{0\}). $$

Note that for $(x,F) \in {}^\theta\widetilde{\mathcal{F}}_k(\mathbf{W},\mathbf{V})$,

$$ \begin{aligned} \dim(\mathbf{F}^\ell_j \cap U_j) &= \dim \mathbf{F}^\ell_j - \delta(j = i, \ell < k), \\ \dim(\mathbf{F}^\ell_j \cap (U^\perp)_j) &= \delta(2m - \ell \ge k, j = \theta(i)). \end{aligned} $$

We have

$$ \dim((F|_{U/U^\perp})^{\ell-1}/(F|_{U/U^\perp})^\ell)_j = \dim(\mathbf{F}^{\ell-1}/\mathbf{F}^\ell)_j - \delta(j = i, \ell = k) - \delta(j = \theta(i), 2m - \ell = k-1). $$

Hence the flag $F|_{U/U^\perp}$ is a flag of type $(\mathbf{i}, \mathbf{a}^{(k)})$. Therefore $(x,F) \mapsto (x|_{U/U^\perp}, F|_{U/U^\perp})$ defines $f_{\mathbf{a}^{(k)}} \colon {}^\theta\widetilde{\mathcal{F}}_k(\mathbf{W},\mathbf{V}) \to {}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a}^{(k)},\Omega}$. We obtain the following diagram:

$$\begin{tikzcd}
{}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a}^{(k)},\Omega} \arrow[d, "{}^\theta\pi_{\mathbf{i},\mathbf{a}^{(k)}}"'] & {}^\theta\widetilde{\mathcal{F}}_k(\mathbf{W},\mathbf{V}) \arrow[l, "f_{\mathbf{a}^{(k)}}"'] \arrow[r, hook] & {}^\theta\widetilde{\mathcal{F}}(\mathbf{W},\mathbf{V}) \arrow[r, hook] \arrow[d] & {}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega} \arrow[d, "{}^\theta\pi_{\mathbf{i},\mathbf{a}}"] \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega} & & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega \arrow[ll, "p"'] \arrow[r, hook, "\iota"] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega}
\end{tikzcd}$$

Claim. *The morphism $f_{\mathbf{a}^{(k)}}$ is an affine bundle of rank $M_k(\mathbf{i}, \mathbf{a}^{(k)})$.*

Proof. Fix $(x_\mathbf{W}, F_\mathbf{W}) \in {}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a}^{(k)},\Omega}$. Note that $(U^\perp)_j = \{0\}$ and $U_j \cong \mathbf{W}_j$ for $j \neq \theta(i)$. If $F \in {}^\theta\mathcal{F}^{(k)}_{\mathbf{i},\mathbf{a};\Omega}$ satisfies $F|_{U/U^\perp} = F_\mathbf{W}$, we have

$$ \mathbf{F}^\ell_i = \mathbf{F}^\ell_{\mathbf{W},i} \ (\ell \ge k), \quad \mathbf{F}^\ell_i = \mathbf{F}^\ell_{\mathbf{W},i} + \mathbf{F}^{k-1}_i \ (\ell < k), \quad \mathbf{F}^\ell_{\theta(i)} = (\mathbf{F}^{2m-k+1}_i)^\perp $$

and $\mathbf{F}^\ell_j = \mathbf{F}^\ell_{\mathbf{W},j}$ $(j \neq i, \theta(i))$. A subspace $\mathbf{F}^{k-1}_i$ is parametrized by a one-dimensional subspace $\mathbf{F}^{k-1}_i/\mathbf{F}^{k-1}_{\mathbf{W},i} \subset \mathbf{V}_i/\mathbf{F}^{k-1}_{\mathbf{W},i}$ such that $\mathbf{F}^{k-1}_i/\mathbf{F}^{k-1}_{\mathbf{W},i} \not\subset U_i/\mathbf{F}^{k-1}_{\mathbf{W},i}$. Hence the fibers of ${}^\theta\mathcal{F}^{(k)}_{\mathbf{i},\mathbf{a},\Omega} \to {}^\theta\mathcal{F}_{\mathbf{i},\mathbf{a}^{(k)},\Omega} \colon F \mapsto F|_{U/U^\perp}$ at $F_\mathbf{W}$ is isomorphic to $\mathbb{A}^{\dim(\mathbf{V}_i/\mathbf{F}^{k-1}_{\mathbf{W},i})-1}$. Note that

$$ \dim(\mathbf{V}_i/\mathbf{F}^{k-1}_{\mathbf{W},i}) - 1 = \sum_{\ell < k, i_\ell = i} a_\ell = \sum_{\ell < k, i_\ell = i} a^{(k)}_\ell. $$

Fix a flag $F \in {}^\theta\mathcal{F}^{(k)}_{\mathbf{i},\mathbf{a};\Omega}$ such that $F|_{U/U^\perp} = F_\mathbf{W}$. Note that $\mathbf{V}_i \supset U_i \cong \mathbf{W}_i$, $\mathbf{V}_{\theta(i)} = U_{\theta(i)}$ and $\mathbf{V}_j = U_j \cong \mathbf{W}_j$ for $j \neq i, \theta(i)$. Assume that $x \in {}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ satisfies the condition that F is x-stable and $x|_{U/U^\perp} = x_\mathbf{W}$.

First, suppose that $h \in \Omega$ satisfies $\mathrm{out}(h) \neq i$ and $\mathrm{in}(h) \neq \theta(i)$. Then x_h coincides with the composition $\mathbf{V}_{\mathrm{out}(h)} \twoheadrightarrow U_{\mathrm{out}(h)}/(U^\perp)_{\mathrm{out}(h)} \cong \mathbf{W}_{\mathrm{out}(h)} \xrightarrow{x_{\mathbf{W},h}} \mathbf{W}_{\mathrm{in}(h)} \cong U_{\mathrm{in}(h)} \subseteq \mathbf{V}_{\mathrm{in}(h)}$. Hence, for such an $h \in \Omega$, x_h is uniquely determined by $x_\mathbf{W}$ and x stabilizes the flag F.

Second, suppose that $h \in \Omega$ satisfies $\operatorname{out}(h) = i$. Take $v \in \mathbf{F}_i^{k-1}$ such that $v \notin U_i$.
If $\operatorname{in}(h) \neq \theta(i)$, x_h is parametrised by $x_h(v) \in \mathbf{F}^{k-1}_{\operatorname{in}(h)}$. Note that

$$\dim \mathbf{F}^{k-1}_{\operatorname{in}(h)} = \sum_{\ell \geq k, i_\ell = \operatorname{in}(h)} a_\ell = \sum_{\ell > k, i_\ell = \operatorname{in}(h)} a_\ell^{(k)},$$

because $\operatorname{in}(h) \neq i, \theta(i)$, $i_k = i$ and $\ell \neq k, 2m-k+1$.
If $\operatorname{out}(h) = i$ and $\operatorname{in}(h) = \theta(i)$, we can regard x_h as a skew-symmetric form on $\mathbf{V}_i$. Since $\mathbf{F}_i^\ell = \mathbf{F}^\ell_{\mathbf{W},i} + \delta(\ell < k)\mathbb{C}v$, the skew-symmetric condition on x is equal to the condition $x(v, \mathbf{F}_i^{2m-k+1} + \mathbb{C}v) = 0$. Then x_h is parametrized by $\left(\mathbf{V}_i/(\mathbf{F}_i^{2m-k+1} + \mathbb{C}v)\right)^*$. Since $v \notin \mathbf{F}_i^{2m-k+1}$ if and only if $2m-k+1 \geq k$, we have

$$\begin{aligned}
&\dim \left(\mathbf{V}_i/(\mathbf{F}_i^{2m-k+1} + \mathbb{C}v)\right)^* = \dim \left(\mathbf{V}/\mathbf{F}_i^{2m-k+1}\right)^* - \delta(2m-k+1 \geq k) \\
= \;&\left(\dim F^{k-1}_{\theta(i)}\right) - \delta(2m-k+1 \geq k) = \left(\sum_{\ell \geq k, i_\ell = \theta(i)} a_\ell\right) - \delta(2m-k+1 \geq k).
\end{aligned}$$

Since $i_k = i \neq \theta(i)$, $i_{2m-k+1} = \theta(i)$, we have $a_\ell = a_\ell^{(k)} + \delta(\ell = 2m-k+1)$ if $i_\ell = \theta(i)$. Thus we obtain

$$\dim \left(\mathbf{V}_i/(\mathbf{F}_i^{2m-k+1} + \mathbb{C}v)\right) = \sum_{\ell > k, i_\ell = \theta(i)} a_\ell^{(k)}.$$

Set

$$\Omega_0 := \{h \in \Omega \mid \operatorname{out}(h) = i, \operatorname{in}(h) = \theta(i)\}, \quad \Omega_1 := \{h \in \Omega \mid \operatorname{out}(h) = i, \operatorname{in}(h) \neq \theta(i)\}.$$

The morphism $\widetilde{\mathcal{F}}_k(\mathbf{W}, \mathbf{V}) \to \{F \in {}^\theta\mathcal{F}^{(k)}_{\mathbf{i},\mathbf{a},\Omega} \mid F|_{U/U^\perp} = F_{\mathbf{W}}\}$ is an affine bundle and its fiber dimension is equal to

$$\begin{aligned}
&\sum_{h \in \Omega_1} \dim(\mathbf{F}^{k-1}_{\operatorname{in}(h)}) + \sum_{h \in \Omega_0} \dim\{\mathbf{V}_i/(\mathbf{F}_i^{2m-k+1} + \mathbb{C}v)\} \\
= &\sum_{h \in \Omega_1, \ell > k, i_\ell \neq \theta(i)} a_\ell^{(k)} + \sum_{h \in \Omega_0, \ell > k, i_\ell = \theta(i)} a_\ell^{(k)} = \sum_{h \in \Omega_0 \sqcup \Omega_1, \ell > k} a_\ell^{(k)}.
\end{aligned}$$

Thus the rank of $f_{\mathbf{a}^{(k)}}$ is equal to

$$\dim(\mathbf{V}_i/\mathbf{F}^{k-1}_{\mathbf{W},i}) - 1 + \sum_{h \in \Omega_0 \sqcup \Omega_1, \ell > k} a_\ell^{(k)} = \sum_{i_\ell = i, \ell < k} a_\ell^{(k)} + \sum_{h \in \Omega_0 \sqcup \Omega_1, k < \ell} a_\ell^{(k)} = M_k(\mathbf{i}, \mathbf{a}^{(k)}).$$

$\square$

By this claim, we have $(f_{\mathbf{a}^{(k)}})_! \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_k(\mathbf{W},\mathbf{V})} = \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a}^{(k)},\Omega}}[-2M_k(\mathbf{i}, \mathbf{a}^{(k)})]$. By Lemma 2.18(2), we obtain

$$\begin{aligned}
\operatorname{Res}_{\mathbf{S}_i,\mathbf{W}}({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) &= ({}^\theta\pi_{\mathbf{i},\mathbf{a};\Omega})_! \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}} = \bigoplus_k ({}^\theta\pi_{\mathbf{i},\mathbf{a}^{(k)}})_! (f_{\mathbf{a}^{(k)}})_! \mathbf{1}_{{}^\theta\widetilde{\mathcal{F}}_k(\mathbf{W},\mathbf{V})} \\
&= \bigoplus_{i_k = i} {}^\theta L_{\mathbf{i},\mathbf{a}^{(k)};\Omega}[-2M_k(\mathbf{i}, \mathbf{a}^{(k)})].
\end{aligned}$$

$\square$

Lemma 4.12. *Let $\mathbf{T}^1$ and $\mathbf{T}^2$ be I-graded vector spaces. Let $\mathbf{W}$ and $\mathbf{V}$ be θ-symmetric I-graded vector spaces such that* $\operatorname{wt} \mathbf{V} = \operatorname{wt} \mathbf{T}^1 + \theta(\operatorname{wt} \mathbf{T}^1) + \operatorname{wt} \mathbf{T}^2 + \theta(\operatorname{wt} \mathbf{T}^2) + \operatorname{wt} \mathbf{W}$. *For $\mathbf{G}_{\mathbf{T}^j}$-equivariant semisimple objects $L_j \in \mathscr{D}(\mathbf{E}_{\mathbf{T}^j,\Omega})$ $(j = 1, 2)$ and a ${}^\theta\mathbf{G}_{\mathbf{W}}$-equivariant semisimple obejct $L \in \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{W},\Omega})$, we have $(L_1 * L_2) * L \cong L_1 * (L_2 * L)$.*

Proof. Let $\mathbf{T}^{12}$ be an I-graded vector space such that $\mathrm{wt}\,\mathbf{T}^{12} = \mathrm{wt}\,\mathbf{T}^1 + \mathrm{wt}\,\mathbf{T}^2$. Let $\mathbf{W}^2$ be a θ-symmetric I-graded vector space such that $\mathrm{wt}\,\mathbf{W}^2 = \mathrm{wt}\,\mathbf{T}^2 + \theta(\mathrm{wt}\,\mathbf{T}^2) + \mathrm{wt}\,\mathbf{W}$.

We denote by $\mathscr{F}$ the variety of pairs (x, F) where $x \in {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}$ and $F = (\mathbf{V} \supset F^1 \supset F^2 \supset F^3 \supset F^4 \supset \{0\}$ is an x-stable flag such that $F^3 = (F^2)^{\perp}, F^4 = (F^1)^{\perp}$, $F^1/F^4 \cong \mathbf{W}^2$ and $F^2/F^3 \cong \mathbf{W}$ as θ-symmetric I-graded vector spaces. Let $\tilde{\mathscr{F}}$ be the variety of pairs $(x, F, \varphi_{\mathbf{W}}, \varphi_{\mathbf{W}^2}, \varphi_1, \varphi_2, \varphi_{\mathbf{T}^2})$ where $(x, F) \in \mathscr{F}$ and $\varphi_{\mathbf{W}^2}\colon F^1/F^4 \cong \mathbf{W}^2$, $\varphi_{\mathbf{W}}\colon F^2/F^3 \cong \mathbf{W}$ as θ-symmetric I-graded vector spaces, and $\varphi_1\colon \mathbf{V}/F^1 \cong \mathbf{T}^1$, $\varphi_2\colon \mathbf{V}/F^2 \cong \mathbf{T}^{12}$ and $\varphi_{\mathbf{T}^2}\colon F^1/F^2 \cong \mathbf{T}^2$ as I-graded vector spaces.

We consider the folowing diagram:

$$\begin{tikzcd}
\mathbf{E}_{\mathbf{T}^1,\Omega} \times \mathbf{E}_{\mathbf{T}^2,\Omega} \times {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} & \tilde{\mathscr{F}} \arrow[l, "u_1"'] \arrow[r, "u_2"] & \mathscr{F} \arrow[r, "u_3"] & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}.
\end{tikzcd}$$

Here $u_1(x, F, \varphi_{\mathbf{W}}, \varphi_{\mathbf{W}^2}, \varphi_1, \varphi_2, \varphi_{\mathbf{T}^2}) = (x^1, x^2, x_{\mathbf{W}})$, where $x_{\mathbf{W}}$, x^1 and x^2 are the restrictions of x to $\mathbf{W}, \mathbf{T}^1$ and $\mathbf{T}^2$ through the isomorphism $\varphi_{\mathbf{W}}, \varphi_1$ and φ_2 respectively, and u_2 and u_3 are natural projections. Note that u_1 is smooth with connected fibers, u_2 is a principal $\mathbf{G}_{\mathbf{T}^1} \times \mathbf{G}_{\mathbf{T}^2} \times {}^{\theta}\mathbf{G}_{\mathbf{W}}$-bundle and u_3 is projective. Then, for $L \in {}^{\theta}\mathscr{P}_{\mathbf{W},\Omega}$, there exists a unique semisimple object $L'' \in \mathscr{D}(\mathscr{F})$ such that $u_1^*(L_1 \boxtimes L_2 \boxtimes L) = u_2^* L''$, we define K by $(u_3)_! L''$. We shall prove $K \cong L_1 * (L_2 * L)$ and $K \cong (L_1 * L_2) * L$.

First, $L_2 * L$ is defined by the following diagram

$$\begin{tikzcd}
\mathbf{E}_{\mathbf{T}^2,\Omega} \times {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} & E_2' \arrow[l, "q_1"'] \arrow[r, "q_2"] & E_2'' \arrow[r, "q_3"] & {}^{\theta}\mathbf{E}_{\mathbf{W}^2,\Omega}.
\end{tikzcd}$$

Here E_2'' is the variety of (y, V) where $y \in {}^{\theta}\mathbf{E}_{\mathbf{W}^2,\Omega}$ and V is an y-stable I-graded vector subspace of $\mathbf{W}^2$ such that $V \supset V^{\perp}$ and $\mathrm{wt}(\mathbf{W}^2/V) = \mathrm{wt}(\mathbf{T}^2)$, and E_2' is the variety of $(y, V, \psi_{\mathbf{W}}, \psi_{\mathbf{T}^2})$ where $(y, V) \in E_2''$ and $\psi_{\mathbf{W}} : V/V^{\perp} \cong \mathbf{W}$ and $\psi_{\mathbf{T}^2} : \mathbf{W}^2/V \cong \mathbf{T}^2$. For $L_2'' \in \mathscr{D}(E_2')$ such that $q_1^*(L_2 \boxtimes L) = q_2^* L_2''$, we have $(q_3)_! L_2'' = L_2 * L$. We consider the diagram

$$\begin{tikzcd}
\mathbf{E}_{\mathbf{T}^1,\Omega} \times \mathbf{E}_{\mathbf{T}^2,\Omega} \times {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} & \mathbf{E}_{\mathbf{T}^1,\Omega} \times E_2' \arrow[l, "\tilde{q}_1"'] \arrow[r, "\tilde{q}_2"] & \mathbf{E}_{\mathbf{T}^1,\Omega} \times E_2'' \arrow[r, "\tilde{q}_3"] & \mathbf{E}_{\mathbf{T}^1,\Omega} \times {}^{\theta}\mathbf{E}_{\mathbf{W}^2,\Omega},
\end{tikzcd}$$

and denote by $L_1''\colon = L_1 \boxtimes L_2'' \in \mathscr{D}(\mathbf{E}_{\mathbf{T}^1,\Omega} \times E_2'')$. Then $\tilde{q}_1^*(L_1 \boxtimes L_2 \boxtimes L) = \tilde{q}_2^* L_1''$ and $(\tilde{q}_3)_! L_1'' = L_1 \boxtimes (L_2 * L)$.

Second, $L_1 * (L_2 * L)$ is defined by the following diagram:

$$\begin{tikzcd}
\mathbf{E}_{\mathbf{T}^1,\Omega} \times {}^{\theta}\mathbf{E}_{\mathbf{W^2},\Omega} & E' \arrow[l, "p_1"'] \arrow[r, "p_2"] & E'' \arrow[r, "p_3"] & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}.
\end{tikzcd}$$

Here E'' is the variety of (y, V) where $y \in {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}$ and V is an y-stable I-graded vector subspace of $\mathbf{V}$ such that $V \supset V^{\perp}$ and $\mathrm{wt}(\mathbf{V}/V) = \mathrm{wt}(\mathbf{T}^1)$, and E' is the variety of $(y, V, \psi_{\mathbf{W}^2}, \psi_{\mathbf{T}^1})$ where $(y, V) \in E''$ and $\psi_{\mathbf{W}^2} : V/V^{\perp} \cong \mathbf{W}^2$ and $\psi_{\mathbf{T}^1} : \mathbf{V}/V \cong \mathbf{T}^1$. For $K'' \in \mathscr{D}(E'')$ such that $p_1^*(L_1 \boxtimes (L_2 * L)) = p_2^* K''$, we have $L_1 * (L_2 * L) = (p_3)_! K''$.

Set $E_1' = \mathbf{E}_{\mathbf{T}^1,\Omega} \times E_2'$, $E_1'' = \mathbf{E}_{\mathbf{T}^1,\Omega} \times E_2''$, $E_{12} = \mathbf{E}_{\mathbf{T}^1,\Omega} \times \mathbf{E}_{\mathbf{T}^2,\Omega} \times {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega}$ and $E_2 = \mathbf{E}_{\mathbf{T}^1,\Omega} \times$

${}^\theta\mathbf{E}_{\mathbf{W}^2,\Omega}$. We consider the following diagram:

$$\begin{tikzcd}
 & & & E_2 & & & \\
 & & E_1'' \arrow[ur, "\tilde q_3"] & \square & E' \arrow[ul, "p_1"'] \arrow[dr, "p_2"] & & \\
 & E_1' \arrow[ur, "\tilde q_2"] \arrow[dl, "\tilde q_1"'] & & \tilde E \arrow[ul, "r_1"'] \arrow[ur, "t_3"] \arrow[dr, "r_2"] & \square & E'' \arrow[dr, "p_3"] & \\
E_{12} & & \tilde{\mathscr{F}} \arrow[ll, "u_1"] \arrow[ul, "v_1"'] \arrow[ur, "t_2"] \arrow[rr, "u_2"'] & & \mathscr{F} \arrow[ur, "s_3"] \arrow[rr, "u_3"'] & & {}^\theta\mathbf{E}_{\mathbf{V},\Omega}
\end{tikzcd}$$

where $\tilde E = \mathscr{F} \times_{E''} E'$. Here $s_3(x, F) = (x, F^1)$, $t_2(x, F, \varphi_{\mathbf{W}}, \varphi_{\mathbf{W}^2}, \varphi_1, \varphi_2) = (x, F, \varphi_{\mathbf{W}^2}, \varphi_1)$, $r_2(x, F, \varphi_{\mathbf{W}^2}, \varphi_1) = (x, F)$ and $t_3(x, F, \varphi_{\mathbf{W}^2}, \varphi_1) = (x, F^1, \varphi_{\mathbf{W}^2}, \varphi_1)$. We define r_1 and v_1 by

$$\begin{aligned} r_1(x, F, \varphi_{\mathbf{W}^2}, \varphi_1) &= (x^1, x_{\mathbf{W}^2}, \varphi_{\mathbf{W}^2}(F_2/F_4)), \\ v_1(x, F, \varphi_{\mathbf{W}}, \varphi_{\mathbf{W}^2}, \varphi_1, \varphi_2, \varphi_{\mathbf{T}^2}) &= (x^1, x_{\mathbf{W}^2}, \varphi_{\mathbf{W}^2}(F_2/F_4), \psi_{\mathbf{W}}, \psi_2), \end{aligned}$$

where $x_{\mathbf{W}^2}, \psi_{\mathbf{W}}$ and ψ_2 are natural morphism induced by using $\varphi_{\mathbf{W}}, \varphi_{\mathbf{W}^2}$ and φ_2.
We have $t_2^* r_1^* L_1'' = v_1^* \tilde q_2^* L_1'' = v_1^* \tilde q_1^* (L_1 \boxtimes L_2 \boxtimes L) = u_1^*(L_1 \boxtimes L_2 \boxtimes L) = u_2^* L'' = t_2^* r_2^* L''$. Since t_2 is a $\mathbf{G}_{\mathbf{T}^2} \times {}^\theta\mathbf{G}_{\mathbf{W}^1}$-principal bundle, we obtain $r_1^* L_1'' = r_2^* L''$. Therefore $p_2^*(s_3)_! L'' = (t_3)_! r_2^* L'' = (t_3)_! r_1^* L'' = p_1^*(q_3)_! L_1'' = p_1^*(L_1 \boxtimes (L_2 * L))$. Thus $(p_3)_!(r_3)_! L'' = L_1 * (L_2 * L)$. We have $K = (u_3)_! L'' = L_1 * (L_2 * L)$.
Similarly, we obtain $K \cong (L_1 * L_2) * L$. Thus the claim follows. □

4.4. **Restriction functor E_i, Induction functors F_i and $F_i^{(a)}$.** We consider the following diagram

$$\mathbf{E}_{\mathbf{T},\Omega} \times {}^\theta\mathbf{E}_{\mathbf{W},\Omega} \xleftarrow{p_1} {}^\theta\mathbf{E}'_\Omega \xrightarrow{p_2} {}^\theta\mathbf{E}''_\Omega \xrightarrow{p_3} {}^\theta\mathbf{E}_{\mathbf{V},\Omega} .$$

Lemma 4.13. *Suppose $\mathbf{T} = \mathbf{S}_i$. Let d_{p_1} and d_{p_2} be the dimension of the fibers of p_1 and p_2, respectively. The we have*

$$d_{p_1} - d_{p_2} = \dim {}^\theta\mathbf{E}''_\Omega - \dim {}^\theta\mathbf{E}_{\mathbf{W},\Omega} = \dim \mathbf{W}_i + \sum_{h \in \Omega:\ \mathrm{out}(h) = i} \dim \mathbf{W}_{\mathrm{in}(h)}.$$

Proof. For a vector space V, we denote by $\mathrm{Alt}(V)$ the set of all skew-symmetric linear maps $V \to V^*$. Let $\mathbf{P}(V)$ denote the projective space of hyperplanes of V. Set $\Omega_0 = \{h \in \Omega \mid \theta(h) = h\}, \Omega_1 = \Omega \backslash \Omega_0$. We have

$$\dim {}^\theta\mathbf{E}_{\mathbf{W},\Omega} = \frac{1}{2} \sum_{h \in \Omega_1} \dim \mathbf{W}_{\mathrm{out}\,(h)} \dim \mathbf{W}_{\mathrm{in}(h)} + \sum_{h \in \Omega_0} \dim \mathrm{Alt}(\mathbf{W}_{\mathrm{out}(h)}).$$

We set

$$\begin{aligned} \Omega_{10} &= \{h \in \Omega_1 \mid \mathrm{out}(h) \neq i, \mathrm{in}(h) \neq i\}, \\ \Omega_{11} &= \{h \in \Omega_1 \mid \mathrm{out}(h) = i\}, \\ \Omega_{12} &= \{h \in \Omega_1 \mid \mathrm{in}(h) = i\}, \\ \Omega_{00} &= \{h \in \Omega_0 \mid (\mathrm{out}(h), \mathrm{in}(h)) = (i, \theta(i)) \text{ or } (\theta(i), i)\}, \\ \Omega_{01} &= \Omega_0 \backslash \Omega_{00}. \end{aligned}$$

Then $\Omega_1 = \Omega_{10} \sqcup \Omega_{11} \sqcup \Omega_{12}$ and $\Omega_0 = \Omega_{00} \sqcup \Omega_{01}$. Note that θ gives bijections $\Omega_{10} \to \Omega_{10}$ and $\Omega_{11} \to \Omega_{12}$. Therefore we have

$$\begin{aligned}\dim {}^{\theta}\mathbf{E}''_{\Omega} &= \dim \mathbf{P}(\mathbf{V}_i) + \frac{1}{2}\sum_{h\in\Omega_{10}} \dim \mathbf{W}_{\mathrm{out}\,(h)} \dim \mathbf{W}_{\mathrm{in}(h)} \\ &\quad + \sum_{h\in\Omega_{11}} \dim \mathbf{V}_i \dim \mathbf{W}_{\mathrm{in}(h)} + \sum_{h\in\Omega_{01}} \dim \mathrm{Alt}(\mathbf{W}_{\mathrm{out}(h)}) \\ &\quad + \sum_{h\in\Omega,\mathrm{out}(h)=i,\mathrm{in}(h)=\theta(i)} \dim \mathrm{Alt}(\mathbf{V}_i) + \sum_{h\in\Omega,\mathrm{out}(h)=\theta(i),\mathrm{in}(h)=i} \dim \mathrm{Alt}(\mathbf{W}_i).\end{aligned}$$

Since $\dim \mathbf{V}_i = \dim \mathbf{W}_i + 1$ and $\dim \mathrm{Alt}(\mathbf{V}_i) - \dim \mathrm{Alt}(\mathbf{W}_i) = \dim \mathbf{W}_i$, we conclude

$$\begin{aligned}&\dim {}^{\theta}\mathbf{E}''_{\Omega} - \dim {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} \\ =\ & \dim \mathbf{W}_i + \sum_{h\in\Omega_{11}} \dim \mathbf{W}_{\mathrm{in}(h)} + \sum_{h\in\Omega,\mathrm{out}(h)=i,\mathrm{in}(h)=\theta(i)} (\dim \mathrm{Alt}(\mathbf{V}_i) - \dim \mathrm{Alt}(\mathbf{W}_i)) \\ =\ & \dim \mathbf{W}_i + \sum_{h\in\Omega,\mathrm{out}(h)=i,\mathrm{in}(h)\neq\theta(i)} \dim \mathbf{W}_{\mathrm{in}(h)} + \sum_{h\in\Omega,\mathrm{out}(h)=i,\mathrm{in}(h)=\theta(i)} \dim \mathbf{W}_i \\ =\ & \dim \mathbf{W}_i + \sum_{h\in\Omega:\ \mathrm{out}(h)=i} \dim \mathbf{W}_{\mathrm{in}(h)}.\end{aligned}$$

□

Definition 4.14.

(i) For $\mathbf{T} = \mathbf{S}_i$ *and a* ${}^{\theta}\mathbf{G}_{\mathbf{W}}$*-equivariant semisimple object* K *in* ${}^{\theta}\mathscr{Q}_{\mathbf{W},\Omega}$*, we define the operator* F_i *by*

$$F_i(K) \colon = (\mathbf{1}_{\mathbf{S}_i} * K)\,[d_{F_i}]$$

where

$$d_{F_i} = d_{p_1} - d_{p_2} = \dim \mathbf{W}_i + \sum_{h\in\Omega:\ \mathrm{out}(h)=i} \dim \mathbf{W}_{\mathrm{in}(h)}.$$

(ii) For $\mathbf{T} = \mathbf{S}_i$*, we define the functor* $E_i \colon \mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}) \to \mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{W},\Omega})$ *by*

$$E_i(K) \colon = \mathrm{Res}_{\mathbf{S}_i,\mathbf{W}}(K)\,[d_{E_i}]$$

where

$$d_{E_i} = d_{F_i} - 2\dim \mathbf{W}_i = -\dim \mathbf{W}_i + \sum_{h\in\Omega:\ \mathrm{out}(h)=i} \dim \mathbf{W}_{\mathrm{in}(h)}.$$

By Prposition 4.11, E_i and F_i induce the restriction functor ${}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega} \to {}^{\theta}\mathscr{Q}_{\mathbf{W},\Omega}$, induction functor ${}^{\theta}\mathscr{Q}_{\mathbf{W},\Omega} \to {}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega}$, respectively.

Definition 4.15. *For* $a \in \mathbb{Z}_{>0}$*, let* $\mathbf{W}$ *and* $\mathbf{V}$ *be* θ*-symmetric* I*-graded vector spaces such that* $\mathrm{wt}(\mathbf{V}) = \mathrm{wt}(\mathbf{W}) + a(\alpha_i + \alpha_{\theta(i)})$*. For a* ${}^{\theta}\mathbf{G}_{\mathbf{W}}$*-equivariant semisimple object* $L \in {}^{\theta}\mathscr{P}_{\mathbf{W},\Omega}$*, we define* $F_i^{(a)}(L) \colon = \mathbf{1}_{\mathbf{S}_i^a} * L[d_a]$ *where*

$$d_a = a\left(\dim \mathbf{W}_i + \sum_{h\in\Omega:\ \mathrm{out}(h)=i} \dim \mathbf{W}_{\mathrm{in}(h)}\right) + \frac{a(a-1)}{2}\sharp\{h \in \Omega|\,\mathrm{out}(h) = i, \mathrm{in}(h) = \theta(i)\}.$$

We call $F_i^{(a)}$ *the* a*-th divided power of* F_i*.*

By Proposition 4.11(1), we have the following lemma.

Lemma 4.16. *The object* ${}^{\theta}L_{\mathbf{i},\mathbf{a};\Omega}$ *is isomorphic to* $F_{i_1}^{(a_1)}F_{i_2}^{(a_2)}\cdots F_{i_m}^{(a_m)}\mathbf{1}_{\mathrm{pt}}$ *up to shift.*

Lemma 4.17. *The operator* $F_i^{(a)}$ *gives a functor* ${}^{\theta}\mathscr{Q}_{\mathbf{W},\Omega} \to {}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega}$ *and satisfy* $F_iF_i^{(a)} = F_i^{(a)}F_i = [a+1]_vF_i^{(a+1)}$.

Proof. By Proposition 4.11(1), $F_i^{(a)}$ gives a functor ${}^{\theta}\mathscr{Q}_{\mathbf{W},\Omega} \to {}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega}$. We have

$$F_iF_i^{(a)}(L) = F_i(\mathbf{1}_{\mathbf{S}_i^a} * L)[d_a] = \mathbf{1}_{\mathbf{S}_i} * (\mathbf{1}_{\mathbf{S}_i^a} * L)v^{d_a+d}$$

where

$$\begin{aligned} d &= \dim \mathbf{W}_i + a + \sum_{h\in\Omega:\ \mathrm{out}(h)=i,\mathrm{in}(h)\neq\theta(i)} \dim \mathbf{W}_{\mathrm{in}(h)} + \sum_{h\in\Omega:\ \mathrm{out}(h)=i,\mathrm{in}(h)=\theta(i)} (\dim \mathbf{W}_{\theta(i)} + a) \\ &= \dim \mathbf{W}_i + a + \sum_{h\in\Omega:\ \mathrm{out}(h)=i} \dim \mathbf{W}_{\mathrm{in}(h)} + a\sharp\{h \in \Omega |\, \mathrm{out}(h) = i, \mathrm{in}(h) = \theta(i)\}. \end{aligned}$$

Note that $\mathbf{1}_{\mathbf{S}_i} * \mathbf{1}_{\mathbf{S}_i^a} = (1 + v^{-2} + \cdots + v^{-2a})\mathbf{1}_{\mathbf{S}_i^{a+1}} = [a+1]_vv^{-a}\mathbf{1}_{\mathbf{S}_i^{a+1}}$ in $\mathbf{E}_{\mathbf{S}_i^{a+1},\Omega}$. By Lemma 4.12, we have

$$\begin{aligned} F_iF_i^{(a)}(L) &= [a+1]_vv^{-a}v^{d_a+d}\mathbf{1}_{\mathbf{S}_i^{a+1}} * L \\ &= [a+1]_vv^{-a}v^{d_a+d-d_{a+1}}F_i^{(a+1)}(L). \end{aligned}$$

Since

$$\begin{aligned} d_a + d &= (a+1)\left(\dim \mathbf{W}_i + \sum_{h\in\Omega:\ \mathrm{out}(h)=i} \dim \mathbf{W}_{\mathrm{in}(h)}\right) + a \\ &\quad + \left(\frac{a(a-1)}{2} + a\right)\sharp\{h \in \Omega |\, \mathrm{out}(h) = i, \mathrm{in}(h) = \theta(i)\} \\ &= d_{a+1} + a, \end{aligned}$$

we conculde $F_iF_i^{(a)} = [a+1]_vF_i^{(a+1)}$. □

4.5. **Commutativity with Fourier transforms.** For two θ-orientations Ω and Ω', we have $\overline{\Omega\backslash\Omega'} = \Omega'\backslash\Omega$. Then we can regard ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega} \to {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}$ and ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'} \to {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}$ as vector bundles and they are the dual vector bundle to each other by the form $\sum_{h\in\Omega\backslash\Omega'} \mathrm{tr}(x_hx_{\bar{h}})$ on ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega} \times {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'}$. We say that $L \in \mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega})$ is $(\mathbb{C}^\times)^{\Omega,\theta}$-monodromic if $H^j(L)$ is locally constant on every $(\mathbb{C}^\times)^{\Omega,\theta}$-orbit on ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}$. Let $\mathscr{D}_{(\mathbb{C}^\times)^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega})$ be the full subcategory of $\mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega})$ consisting of $(\mathbb{C}^\times)^{\Omega,\theta}$-monodromic objects. Hence we have the Fourier transform $\Phi_{\mathbf{V}}^{\Omega\Omega'}\colon \mathscr{D}_{(\mathbb{C}^\times)^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}) \to \mathscr{D}_{(\mathbb{C}^\times)^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'})$. The following lemma is obvious.

Lemma 4.18. *For three θ-orientations Ω, Ω' and Ω'', we have*

$$\Phi_{\mathbf{V}}^{\Omega'\Omega''} \circ \Phi_{\mathbf{V}}^{\Omega\Omega'} \cong a^* \circ \Phi_{\mathbf{V}}^{\Omega\Omega''} : \mathscr{D}_{(\mathbb{C}^\times)^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}) \to \mathscr{D}_{(\mathbb{C}^\times)^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega''})$$

where $a : {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega''} \to {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega''}$ *is defined by* $x_h \mapsto -x_h$ *or* x_h *according that* $h \in \Omega'' \cap \overline{\Omega'} \cap \Omega$ *or not. In particular,* $\mathscr{D}_{(\mathbb{C}^\times)^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega})$ *does not depend on* Ω.

Since any object in ${}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega}$ is ${}^{\theta}\mathbf{G}_{\mathbf{V}} \times (\mathbb{C}^\times)^{\Omega,\theta}$-equivariant, it is a monodromic object. By the commutativity between E_i, F_i and $(\mathbb{C}^\times)^{\Omega,\theta}$-action, the functors E_i and F_i preserve the category $(\mathbb{C}^\times)^{\Omega,\theta}$-monodromic objects.

Theorem 4.19. *Let* $\mathbf{V}$ *and* $\mathbf{W}$ *be* θ*-symmetric* I*-graded vector spaces such that* $\operatorname{wt}\mathbf{V} = \operatorname{wt}\mathbf{W} + \alpha_i + \alpha_{\theta(i)}$*, and* Ω *and* Ω' *be two* θ*-symmetric orientations.*

(1) *Let* F_i^{Ω} *and* $F_i^{\Omega'}$ *be the induction functors with respect to* Ω *and* Ω'*, respectively. For a* ${}^{\theta}\mathbf{G}_{\mathbf{W}}$*-equivariant semisimple obejct* $L \in {}^{\theta}\mathscr{Q}_{\mathbf{W},\Omega}$*, we have* $\Phi_{\mathbf{V}}^{\Omega\Omega'} \circ F_i^{\Omega}(L) \cong F_i^{\Omega'} \circ \Phi_{\mathbf{W}}^{\Omega\Omega'}(L)$*.*

(2) *Let* ${E_i}^{\Omega}$ *and* ${E_i}^{\Omega'}$ *be the restriction functors with respect to* Ω *and* Ω'*, respectively. For a* ${}^{\theta}\mathbf{G}_{\mathbf{V}}$*-equivariant semisimple obejct* $K \in {}^{\theta}\mathscr{Q}_{\mathbf{W},\Omega}$*, we have* $\Phi_{\mathbf{W}}^{\Omega\Omega'} \circ {E_i}^{\Omega}(K) \cong {E_i}^{\Omega'} \circ \Phi_{\mathbf{V}}^{\Omega\Omega'}(K)$*.*

(3) *The Fourier transform* $\Phi_{\mathbf{V}}^{\Omega\Omega'}$ *gives an isomorphism between* ${}^{\theta}\mathscr{P}_{\mathbf{V},\Omega}$ *and* ${}^{\theta}\mathscr{P}_{\mathbf{V},\Omega'}$ *and an equivalence between* ${}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega}$ *and* ${}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega'}$*.*

Proof. (1) Let us define the fibre products $E_1, E_2, E_3, E_1', E_2'$ and E_3' by

$$\begin{aligned}
E_1 &\colon = {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} \times_{{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}'_{\Omega\cap\Omega'}, & E_1' &\colon = {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega'} \times_{{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}'_{\Omega\cap\Omega'}, \\
E_2 &\colon = {}^{\theta}\mathbf{E}'_{\Omega\cap\Omega'} \times_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}, & E_2' &\colon = {}^{\theta}\mathbf{E}'_{\Omega\cap\Omega'} \times_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}, \\
E_3 &\colon = {}^{\theta}\mathbf{E}''_{\Omega\cap\Omega'} \times_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}, & E_3' &\colon = {}^{\theta}\mathbf{E}''_{\Omega\cap\Omega'} \times_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'}.
\end{aligned}$$

Note that E_1' and E_2' are the dual vector bundle of E_1 and E_2 over ${}^{\theta}\mathbf{E}'_{\Omega\cap\Omega'}$ respectively, and E_3' is the dual vector bundle of E_3 over ${}^{\theta}\mathbf{E}''_{\Omega\cap\Omega'}$. We denote by $\Phi_{E_j} \colon \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}(E_j) \to \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}(E_j')$ $(j = 1, 2, 3)$ and $\Phi' \colon \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}'_{\Omega}) \to \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}{\mathbf{E}'_{\Omega}}^{*})$ the Fourier transforms. For simplicity, we denote by $\Phi_{\mathbf{V}}, \Phi_{\mathbf{W}}$ insted of $\Phi_{\mathbf{V}}^{\Omega\Omega'}, \Phi_{\mathbf{W}}^{\Omega\Omega'}$, respectively.

We denote by u_1 and u_1' the projections $E_1 \to {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega}$ and $E_1' \to {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega'}$, respectively. Let $\tilde{p_1}, \tilde{p_1'}, \iota_2$ and ι_2' be the canonical maps ${}^{\theta}\mathbf{E}'_{\Omega} \to E_1, {}^{\theta}\mathbf{E}'_{\Omega'} \to E_1', {}^{\theta}\mathbf{E}'_{\Omega} \to E_2$ and ${}^{\theta}\mathbf{E}'_{\Omega'} \to E_2'$, respectively. Then we obtain the following Cartesian diagram of the vector bundles on ${}^{\theta}\mathbf{E}'_{\Omega\cap\Omega'}$:

$$\begin{tikzcd}
{}^{\theta}\mathbf{E}'_{\Omega'} \arrow[r, "\iota_2'"] \arrow[d, "\tilde{p_1'}"'] \arrow[dr, phantom, "\square"] & E_2' \arrow[d, "{}^{t}\iota_2"] \\
E_1' \arrow[r, "{}^{t}\tilde{p_1}"] & ({}^{\theta}\mathbf{E}'_{\Omega})^{*}
\end{tikzcd}$$

Moreover let u_3 and u_3' be the projections $E_3 \to {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}$ and $E_3' \to {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'}$, respectively, $\tilde{p_2}, \tilde{p_2'}, \iota_3$ and ι_3' the canonical maps ${}^{\theta}\mathbf{E}''_{\omega} \to E_3$ and ${}^{\theta}\mathbf{E}''_{\Omega'} \to E_3'$, repectively. We obtain the following diagram:

$$\begin{tikzcd}
E_1 \arrow[d, "u_1"'] & E_2 \arrow[r, "\widetilde{p_2}"] \arrow[dr, phantom, "\square"] & E_3 \arrow[dr, "u_3"] & \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega} \arrow[d] & {}^\theta\mathbf{E}'_\Omega \arrow[l, "p_1"'] \arrow[r, "p_2"] \arrow[ul, "\widetilde{p_1}"'] \arrow[u, "\iota_2"'] \arrow[d] & {}^\theta\mathbf{E}''_\Omega \arrow[r, "p_3"] \arrow[u, "\iota_3"'] \arrow[d] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \arrow[d] \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'} & {}^\theta\mathbf{E}'_{\Omega\cap\Omega'} \arrow[l] \arrow[r] & {}^\theta\mathbf{E}''_{\Omega\cap\Omega'} \arrow[r] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'} \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega'} \arrow[u] & {}^\theta\mathbf{E}'_{\Omega'} \arrow[l, "p'_1"'] \arrow[r, "p'_2"] \arrow[u] \arrow[dl, "\widetilde{p'_1}"'] \arrow[d, "\iota'_2"'] \arrow[dr, phantom, "\square"] & {}^\theta\mathbf{E}''_{\Omega'} \arrow[r, "p'_3"] \arrow[u] \arrow[d, "\iota'_3"'] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega'} \arrow[u] \\
E'_1 \arrow[u, "u'_1"'] \arrow[d, "{}^t\widetilde{p_1}"'] \arrow[r, phantom, "\square"] & E'_2 \arrow[r, "\widetilde{p'_2}"'] \arrow[dl, "{}^t\iota_2"'] & E'_3 \arrow[ur, "u'_3"'] & \\
({}^\theta\mathbf{E}'_\Omega)^* & & &
\end{tikzcd}$$

Let L be ${}^\theta\mathbf{G}_\mathbf{W}$-equivariant semisimple complex on ${}^\theta\mathbf{E}_{\mathbf{W},\Omega}$, L'' a unique semisimple complex such that $p_2^*L'' = p_1^*L$, and K'' a unique semisimple complex such that ${p'_1}^*\Phi_\mathbf{W}(L) = {p'_2}^*K''$.
By Proposition 2.21, we have

$$\begin{aligned}
\widetilde{p'_2}^*\Phi_{E_3}((\iota_3)_!L'') &= \Phi_{E_2}(\widetilde{p_2}^*(\iota_3)_!L'') = \Phi_{E_2}((\iota_2)_!p_2^*L'') = \Phi_{E_2}((\iota_2)_!p_1^*L) \\
&= {}^t\iota_2^*\Phi'(\widetilde{p_1}^*u_1^*L)[d_2] \\
&= {}^t\iota_2^*({}^tp_1)_!(u'_1)^*\Phi_\mathbf{W}(L)[d_2+d_1] \\
&= (\iota'_2)_!\widetilde{p'_1}^*\Phi_\mathbf{W}(L)[d_1+d_2] = (\iota'_2)_!(p'_1)^*\Phi_\mathbf{W}(L)[d_1+d_2] \\
&= (\iota'_2)_!(p'_2)^*K''[d_1+d_2] = \widetilde{p'_2}^*(\iota'_3)_!K''[d_1+d_2]
\end{aligned}$$

where

$$d_1 = \mathrm{rank}(E_1) - \mathrm{rank}({}^\theta\mathbf{E}'_\Omega), \quad d_2 = \mathrm{rank}(E_2) - \mathrm{rank}({}^\theta\mathbf{E}'_\Omega).$$

Hence $\Phi_{E_3}((\iota_3)_!L) = (\iota'_3)_!K''[d_1+d_2]$. Then

$$\Phi_\mathbf{V}((p_3)_!L'') = (u'_3)_!\Phi_{E_3}((\iota_3)_!L'') = (u'_3)_!(\iota'_3)_!K''[d_1+d_2] = (p'_3)_!K''[d_1+d_2].$$

We have

$$\Phi_\mathbf{V}\circ F_i^\Omega(L) = F_i^{\Omega'}\circ\Phi_\mathbf{W}(L)[d]$$

where

$$d = d_1 + d_2 + \sum_{i\xrightarrow{\Omega}\xi}\dim W_\xi - \sum_{i\xrightarrow{\Omega'}\eta}\dim W_\eta.$$

Now we suppose $\Omega\backslash\Omega' = \{h, \theta(h)\}$ and put $\mathrm{out}(h) = k, \mathrm{in}(h) = \ell$. When $k = i$, we have $\sum_{i\xrightarrow{\Omega}\xi}\dim W_\xi - \sum_{i\xrightarrow{\Omega'}\eta}\dim W_\eta = \dim \mathbf{W}_\ell$. If $\ell \neq \theta(i)$, we have $d_2 = 0$ and $d_1 = \dim\mathbf{W}_i\dim\mathbf{W}_\ell - \dim\mathbf{V}_i\dim\mathbf{V}_\ell = -\dim\mathbf{W}_\ell$. If $\ell = \theta(i)$, we have $d_2 = 0$ and $d_1 = \dim\mathrm{Alt}(\mathbf{W}_i, \mathbf{W}_{\theta(i)}) - \dim\mathrm{Alt}(\mathbf{V}_i, \mathbf{V}_{\theta(i)}) = -\dim\mathbf{W}_{\theta(i)}$. Thus we obtain $\Phi_\mathbf{V}\circ F_i^\Omega(L) = F_i^{\Omega'}\circ\Phi_\mathbf{W}(L)$. When $k = \theta(i)$, we can prove the claim by the same way.

(2) We may suppose $\Omega\backslash\Omega' = \{h, \theta(h)\}$ and put $\mathrm{out}(h) = k, \mathrm{in}(h) = \ell$.
We considet the following diagram:

$$\begin{tikzcd}
{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} \arrow[d] & {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega} \arrow[l, "p"'] \arrow[r, "\iota"] \arrow[d] & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega} \arrow[d] \\
{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'} & {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'} \arrow[l] \arrow[r] & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'} \\
{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega'} \arrow[u] & {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega'} \arrow[l, "p'"'] \arrow[r, "\iota'"] \arrow[u] & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'} \arrow[u]
\end{tikzcd}$$

If $k, \ell \neq i, \theta(i)$, the above four diagrams are cartesian. Then the commutativity is clear.
When $k = i$, we consider the two fiber products by the following:

$$\begin{tikzcd}
{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} \arrow[ddd] & & {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega} \arrow[ll, "p"'] \arrow[rr, "\iota"] \arrow[dl, "q_2"'] \arrow[ddd] & & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega} \arrow[ddd] \\
& E \arrow[ul, "q_1"] \arrow[ddr] & & \square & \\
& \square & & & \\
{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'} & & {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'} \arrow[ll] \arrow[rr] & & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'} \\
& & & \square & \\
& \square & & E' \arrow[uul] \arrow[dr, "r_1"] & \\
{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega'} \arrow[uuu] & & {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega'} \arrow[ll, "p'"'] \arrow[rr, "\iota'"] \arrow[uuu] \arrow[ur, "r_2"] & & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'} \arrow[uuu]
\end{tikzcd}$$

where

$$E\colon = {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} \times_{{}^{\theta}\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'}, \quad E'\colon = {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega'} \times_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}} {}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'}.$$

We can regard E and E' as the dual vector bundle of ${}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega'}$ and ${}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega}$ on ${}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'}$ respectively. We can regard r_2 as the transpose of q_2. We denote by Φ and Φ' the Fourier transforms

$$\begin{aligned}
\Phi &: \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}(E) \to \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega'}), \\
\Phi' &: \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}({}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega}) \to \mathscr{D}_{(\mathbb{C}^{\times})^{\Omega,\theta}-\mathrm{mono}}(E').
\end{aligned}$$

Then, for $K \in {}^{\theta}\mathscr{Q}_{\mathbf{V},\Omega}$ we have

$$\Phi_{\mathbf{W}}(p_!\iota^* K) = p'_!\Phi((q_2)_!\iota^* K) = p'_! r_2^* \Phi'(\iota^* K)[d] = p'_! r_2^* r_1^* \Phi_{\mathbf{V}}(K)[d] = p'_!(\iota')^*\Phi_{\mathbf{V}}(K)[d],$$

where $d = \mathrm{rank}(E) - \mathrm{rank}({}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega})$. If $\ell \neq \theta(i)$, we have $\mathrm{rank}(E) = \dim \mathbf{W}_i \dim \mathbf{W}_\ell$ and $\mathrm{rank}({}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega}) = \dim \mathbf{V}_i \dim \mathbf{V}_\ell$. Since $\mathbf{V}_\ell = \mathbf{W}_\ell$, we have $d = -\dim \mathbf{W}_\ell$. If $\ell = \theta(i)$, we have $\mathrm{rank}(E) = \dim \mathrm{Alt}(\mathbf{W}_i, \mathbf{W}_{\theta(i)})$ and $\mathrm{rank}({}^{\theta}\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega}) = \dim \mathrm{Alt}(\mathbf{V}_i, \mathbf{V}_{\theta(i)})$. Then $d = -\dim \mathbf{W}_{\theta(i)}$. Since $\Omega\backslash\Omega' = \{i \to \ell, \theta(\ell) \to \theta(i)\}$, we have $d_{E_i^{\Omega}} - \dim \mathbf{W}_\ell = d_{E_i^{\Omega'}}$. Thus $\Phi_{\mathbf{W}} \circ E_i^{\Omega}(K) = E_i^{\Omega'} \circ \Phi_{\mathbf{V}}$.

When $k = \theta(i)$, we obtain the following diagram:

$$\begin{tikzcd}
{}^\theta\mathbf{E}_{\mathbf{W},\Omega} \arrow[ddd] & & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega \arrow[ll, "p"'] \arrow[rrr, "\iota"] \arrow[dr, "\iota_2"] \arrow[ddd] & & & {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \arrow[ddd] \\
& \square & & F \arrow[urr] \arrow[ddl] & & \\
& & & & \square & \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'} & & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'} \arrow[ll] \arrow[rrr] & & & {}^\theta\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'} \\
& \square & & & & \\
& F' \arrow[uur] \arrow[dl] & & \square & & \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega'} \arrow[uuu] & & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega'} \arrow[ll, "p'"'] \arrow[rrr, "\iota'"] \arrow[ul, "p'_2"'] \arrow[uuu] & & & {}^\theta\mathbf{E}_{\mathbf{V},\Omega'} \arrow[uuu]
\end{tikzcd}$$

Here

$$F\colon = {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \times_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega\cap\Omega'}} {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'}, \quad F'\colon = {}^\theta\mathbf{E}_{\mathbf{W},\Omega'} \times_{{}^\theta\mathbf{E}_{\mathbf{W},\Omega\cap\Omega'}} {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_{\Omega\cap\Omega'}.$$

We regard p'_2 as the transpose of ι_2. Hence we can prove the claim by the similar way.
(3) The claim follows from Proposition 2.21(2) and the commutativity of F_i and $\Phi_{\mathbf{V}}^{\Omega\Omega'}$. $\square$

Similarly, we can prove the commutativity of $F_i^{(a)}$'s and the Fourier transforms. We omit the proof.

Proposition 4.20. *Let* $\mathbf{W}$ *and* $\mathbf{V}$ *be θ-symmetric I-graded vector spaces such that* $\mathrm{wt}(\mathbf{V}) = \mathrm{wt}(\mathbf{W}) + a(\alpha_i + \alpha_{\theta(i)})$. *Let* ${F_i^{(a)}}^{\Omega}$ *and* ${F_i^{(a)}}^{\Omega'}$ *be the a-th divided powers with respect to θ-orientations Ω and Ω', respectively. For a ${}^\theta\mathbf{G}_{\mathbf{W}}$-equivariant semisimple obejct $L \in {}^\theta\mathscr{Q}_{\mathbf{W},\Omega}$, we have* $\Phi_{\mathbf{V}}^{\Omega\Omega'} \circ {F_i^{(a)}}^{\Omega}(L) \cong {F_i^{(a)}}^{\Omega'} \circ \Phi_{\mathbf{W}}^{\Omega\Omega'}(L)$.

5. A GEOMETRIC CONSTUCTION OF SYMMETRIC CRYSTALS

5.1. **Grothendieck group.** For a θ-orientation Ω and a θ-symmetric and I-graded vector space $\mathbf{V}$, we define ${}^\theta K_{\mathbf{V},\Omega}$ as the Grothendieck group of ${}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$. Namely ${}^\theta K_{\mathbf{V},\Omega}$ is generated by (L) for $L \in {}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$ with the relation $(L) = (L') + (L'')$ when $L \cong L' \oplus L''$. This is a $\mathbb{Z}[v, v^{-1}]$-module by $v(L) = (L[1])$ and $v^{-1}(L) = (L[-1])$ for $L \in {}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$. Hence, ${}^\theta K_{\mathbf{V},\Omega}$ is a free $\mathbb{Z}[v, v^{-1}]$-module with a basis $\{(L) \mid L \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}\}$. For another θ-symmetric and I-graded vector space $\mathbf{V}'$ such that $\mathrm{wt}\,\mathbf{V} = \mathrm{wt}\,\mathbf{V}'$, we have ${}^\theta K_{\mathbf{V},\Omega} \cong {}^\theta K_{\mathbf{V}',\Omega}$. We define

$${}^\theta K_\Omega\colon = \bigoplus_{\mathbf{V}} {}^\theta K_{\mathbf{V},\Omega}$$

where $\mathbf{V}$ runs over the isomorphism classes of θ-symmetric I-graded vector spaces. For two θ-orientations Ω and Ω', the Fourier transform induces an equivalence ${}^\theta\mathscr{Q}_{\mathbf{V},\Omega} \to {}^\theta\mathscr{Q}_{\mathbf{V},\Omega'}$ and the isomorphism ${}^\theta K_{\mathbf{V},\Omega} \xrightarrow{\sim} {}^\theta K_{\mathbf{V},\Omega'}$. Therefore ${}^\theta K_\Omega \cong {}^\theta K_{\Omega'}$.

We set ${}^\theta K = {}^\theta K_\Omega, {}^\theta\mathscr{P}_{\mathbf{V}} = {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$. By Lemma 4.18, they are well-defined.

5.2. **Actions of E_i and F_i.** The functors E_i and $F_i^{(a)}$ induce the action on ${}^\theta K_\Omega$.Since E_i and F_i commute with the Fourier transforms, they also act on ${}^\theta K$. The submodule ${}^\theta K'\colon = \sum_{(\mathbf{i},\mathbf{a})} \mathbb{Z}[v, v^{-1}]({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) \subset {}^\theta K$ is stable by E_i and F_i by Proposition 4.11. We define

$$T_i|_{{}^\theta K_{\mathbf{V},\Omega}} = v^{-(\alpha_i, \mathrm{wt}\,\mathbf{V})}\,\mathrm{id}_{{}^\theta K_{\mathbf{V},\Omega}}.$$

Proposition 5.1. *The operators* E_i, F_i *and* T_i $(i \in I)$ *regarded as operators on* ${}^{\theta}K'$ *satisfy*

$$E_iF_j - v^{-(\alpha_i,\alpha_j)}F_jE_i = \delta_{ij} + \delta_{\theta(i),j}T_i$$

and

$$T_iE_jT_i^{-1} = v^{(\alpha_i+\alpha_{\theta(i)},\alpha_j)}E_j, \quad T_iF_jT_i^{-1} = v^{(\alpha_i+\alpha_{\theta(i)},-\alpha_j)}F_j.$$

Proof. We take θ-symmetric I-graded vector spaces $\mathbf{W}, \mathbf{V}, \mathbf{U}$ and $\mathbf{X}$ such that $\mathrm{wt}(\mathbf{V}) = \mathrm{wt}(\mathbf{W})+\alpha_j+\alpha_{\theta(j)}$, $\mathrm{wt}(\mathbf{U}) = \mathrm{wt}(\mathbf{W})+\alpha_i+\alpha_{\theta(i)}$ and $\mathrm{wt}(\mathbf{X}) = \mathrm{wt}(\mathbf{W})+\alpha_j+\alpha_{\theta(j)}+\alpha_i+\alpha_{\theta(i)}$. We consider the following diagram:

$$\begin{tikzcd}
\mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{W},\Omega}) \arrow[r, "F_j"] \arrow[d, "E_i"'] & \mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}) \arrow[d, "E_i"] \\
\mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{U},\Omega}) \arrow[r, "F_j"'] & \mathscr{D}({}^{\theta}\mathbf{E}_{\mathbf{X},\Omega})
\end{tikzcd}$$

First, we have

$$E_iF_j{}^{\theta}L_{\mathbf{i},\mathbf{a};\Omega} \;=\; \delta_{ij}{}^{\theta}L_{(i,\mathbf{i},\theta(i)),(0,\mathbf{a},0);\Omega}[c_{\mathbf{a}}] \oplus \delta_{\theta(i),j}{}^{\theta}L_{(\theta(i),\mathbf{i},i),(0,\mathbf{a},0);\Omega}[c_{\mathbf{a}_\theta}] \oplus \bigoplus_{\mathbf{a}'}{}^{\theta}L_{(j,\mathbf{i},\theta(j)),(1,\mathbf{a}',1);\Omega}[c_{\mathbf{a}'}],$$

where

$$\begin{aligned}
c_{\mathbf{a}} &= \dim\mathbf{W}_i + \sum_{i\to\eta}\dim\mathbf{W}_\eta - \dim\mathbf{X}_i + \sum_{i\to\xi}\dim\mathbf{X}_\xi - 2M_1((i,\mathbf{i},\theta(i)),(0,\mathbf{a},0)),\\
c_{\mathbf{a}_\theta} &= \dim\mathbf{W}_{\theta(i)} + \sum_{\theta(i)\to\eta}\dim\mathbf{W}_\eta - \dim\mathbf{X}_i + \sum_{i\to\xi}\dim\mathbf{X}_\xi - 2M_{2m+1}((\theta(i),\mathbf{i},i),(0,\mathbf{a},0)),\\
c_{\mathbf{a}'} &= \dim\mathbf{W}_j + \sum_{j\to\eta}\dim\mathbf{W}_j - \dim\mathbf{X}_i + \sum_{i\to\xi}\dim\mathbf{X}_\xi - 2M_{k+1}((j,\mathbf{i},\theta(j)),(1,\mathbf{a}',1)).
\end{aligned}$$

Here $\mathbf{a}'$ runs over the sequences $\mathbf{a}^{(k)}$ $(1 \le k \le m, i_k = i, \theta(i))$.
If $i = j$, we have $c_{\mathbf{a}} = 0$ by $\mathbf{W} = \mathbf{X}$ and

$$M_1((i,\mathbf{i},\theta(i)),(0,\mathbf{a},0)) = \sum_{i\to i_\ell} a_\ell = \sum_{i\to\eta}\dim\mathbf{W}_\eta.$$

If $\theta(i) = j$, we have

$$c_{\mathbf{a}_\theta} = \sum_{\eta\to i}\dim\mathbf{W}_\eta + \sum_{i\to\eta}\dim\mathbf{W}_\eta - 2\dim\mathbf{W}_i = -(\alpha_i, \mathrm{wt}(\mathbf{V}))$$

by $\mathbf{W} = \mathbf{X}$,

$$M_{2m+1}((\theta(i),\mathbf{i},i),(0,\mathbf{a},0)) = \sum_{i_\ell=i} a_\ell = \dim\mathbf{W}_i$$

and $\sum_{\theta(i)\to\eta}\dim\mathbf{W}_\eta = \sum_{\eta\to i}\dim\mathbf{W}_\eta$.
On the other hand, we have

$$F_jE_i{}^{\theta}L_{\mathbf{i},\mathbf{a};\Omega} = \bigoplus_{\mathbf{a}'}{}^{\theta}L_{(j,\mathbf{i},\theta(j)),(1,\mathbf{a}',1);\Omega}[d_{\mathbf{a}'}],$$

where

$$d_{\mathbf{a}'} = -\dim\mathbf{U}_i + \sum_{i\to\xi}\dim\mathbf{U}_\xi + \dim\mathbf{U}_j + \sum_{j\to\eta}\dim\mathbf{U}_\eta - 2M_k(\mathbf{i},\mathbf{a}').$$

and $\mathbf{a}'$ runs over the sequences $\mathbf{a}^{(k)}$ $(1 \leq k \leq m, i_k = i, \theta(i))$.
We have

$$M_{k+1}((j, \mathbf{i}, \theta(j)), (1, \mathbf{a}', 1)) - M_k(\mathbf{i}, \mathbf{a}') = \begin{cases} 1 + \sharp\{i \xrightarrow{\Omega} \theta(i)\} & (j = i) \\ 0 & (j = \theta(i)) \\ \sharp\{i \xrightarrow{\Omega} \theta(j)\} & (j \neq i, \theta(i)) \end{cases}$$

and

$$\begin{aligned} &\left(\dim \mathbf{W}_j + \sum_{j \to \eta} \dim \mathbf{W}_j - \dim \mathbf{X}_i + \sum_{i \to \xi} \dim \mathbf{X}_\xi\right) \\ &\quad - \left(- \dim \mathbf{U}_i + \sum_{i \to \xi} \dim \mathbf{U}_\xi + \dim \mathbf{U}_j + \sum_{j \to \eta} \dim \mathbf{U}_\eta\right) \\ &= \begin{cases} 2\sharp\{i \xrightarrow{\Omega} \theta(i)\} & (j = i) \\ \sharp\{i \xrightarrow{\Omega} \theta(i)\} + \sharp\{\theta(i) \xrightarrow{\Omega} i\} & (j = \theta(i)) \\ \sharp\{i \xrightarrow{\Omega} j\} + \sharp\{j \xrightarrow{\Omega} i\} + 2\sharp\{i \xrightarrow{\Omega} \theta(j)\} & (j \neq i, \theta(i)) \end{cases} . \end{aligned}$$

Thus $c_{\mathbf{a}'} - d_{\mathbf{a}'} = -(\alpha_i, \alpha_j)$. We conclude

$$E_i F_j({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) - v^{-(\alpha_i, \alpha_j)} F_j E_i({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) = \delta_{ij}({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) + \delta_{\theta(i),j} T_i({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}).$$

The relations $T_i E_j T_i^{-1} = v^{(\alpha_i + \alpha_{\theta(i)}, \alpha_j)} E_j$ and $T_i F_j T_i^{-1} = v^{(\alpha_i + \alpha_{\theta(i)}, -\alpha_j)} F_j$ are obvious. $\square$

5.3. **Key estimates of coefficients.** Let Ω be a θ-orientation and suppose that a vertex i is a sink. For a θ-symmetric I-graded vector space $\mathbf{V}$ and $r \in \mathbb{Z}_{\geq 0}$, we define

$${}^\theta\mathbf{E}_{\mathbf{V},\Omega,r} \colon = \left\{ x \in {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \;\middle|\; \dim \operatorname{Coker}\left(\bigoplus_{h \in \Omega; \mathrm{in}(h) = i} \mathbf{V}_{\mathrm{out}(h)} \to \mathbf{V}_i \right) = r \right\}.$$

Then we have ${}^\theta\mathbf{E}_{\mathbf{V},\Omega} = \sqcup_{r \geq 0} {}^\theta\mathbf{E}_{\mathbf{V},\Omega,r}$, and ${}^\theta\mathbf{E}_{\mathbf{V},\Omega,\geq r} \colon = \sqcup_{r' \geq r} {}^\theta\mathbf{E}_{\mathbf{V},\Omega,r'}$ is a closed subset of ${}^\theta\mathbf{E}_{\mathbf{V},\Omega}$.

Definition 5.2. *For $L \in {}^\theta\mathscr{P}_{\mathbf{V}}$ and $i \in I$, choose a θ-orientation Ω such that i is a sink with respect to Ω, and regard L as an element of ${}^\theta\mathscr{P}_{\mathbf{V},\Omega}$. We define $\varepsilon_i(L)$ as the largest integer r satisfying* $\operatorname{Supp}(L) \subset {}^\theta\mathbf{E}_{\mathbf{V},\Omega,\geq r}$. *This does not depend on the choice of Ω.*

Note that $0 \leq \varepsilon_i(L) \leq \dim V_i$.
We shall prove the following key estimates with respect to $F_i(L)$ and $E_i(L)$.

Theorem 5.3. *Assume that θ-symmetric and I-graded vector spaces $\mathbf{V}$ and $\mathbf{W}$ satisfy* $\operatorname{wt} \mathbf{V} = \operatorname{wt} \mathbf{W} + \alpha_i + \alpha_{\theta(i)}$. *Fix a θ-orientation Ω such that the vertex i is a sink.*

(1) *For $L \in {}^\theta\mathscr{P}_{\mathbf{W},\Omega}$, there exists a unique simple perverse sheaf $L_0 \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$ such that $\varepsilon_i(L_0) = \varepsilon_i(L) + 1$ and*

$$F_i(L) = [\varepsilon_i(L) + 1]_v (L_0) + \sum_{L' \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega} \colon \varepsilon_i(L') > \varepsilon_i(L) + 1} a_{L'}(L')$$

for $a_{L'} \in v^{2 - \varepsilon_i(L')}\mathbb{Z}[v]$.
We define the map $\tilde{F}_i \colon {}^\theta\mathscr{P}_{\mathbf{W}} \cong {}^\theta\mathscr{P}_{\mathbf{W},\Omega} \to {}^\theta\mathscr{P}_{\mathbf{V},\Omega} \cong {}^\theta\mathscr{P}_{\mathbf{V}}$ by $\tilde{F}_i(L) = L_0$. It does not depend on the choice of Ω.

(2) *Let* $K \in {}^{\theta}\mathscr{P}_{\mathbf{V},\Omega}$*. If* $\varepsilon_i(K) > 0$*, there exists a unique simple perverse sheaf* $K_0 \in {}^{\theta}\mathscr{P}_{\mathbf{W},\Omega}$ *such that* $\varepsilon_i(K_0) = \varepsilon_i(K) - 1$ *and*

$$E_i(K) = v^{1-\varepsilon_i(K)}(K_0) + \sum_{K' \in {}^{\theta}\mathscr{P}_{\mathbf{W},\Omega}\colon \varepsilon_i(K') > \varepsilon_i(K)-1} b_{K'}(K')$$

for $b_{K'} \in v^{-\varepsilon_i(K')+1}\mathbb{Z}[v]$*. Here we regard* $K_0 = 0$ *if* $\varepsilon_i(K) = 0$*.*
We define the map $\tilde{E}_i \colon {}^{\theta}\mathscr{P}_{\mathbf{V}} \cong {}^{\theta}\mathscr{P}_{\mathbf{V},\Omega} \to {}^{\theta}\mathscr{P}_{\mathbf{W},\Omega} \sqcup \{0\} \cong {}^{\theta}\mathscr{P}_{\mathbf{W}} \sqcup \{0\}$ *by* $\tilde{E}_i(K) = K_0$ *if* $\varepsilon_i(K) > 0$ *and* $\tilde{E}_i(K) = 0$ *if* $\varepsilon_i(K) = 0$*. It does not depend on the choise of* Ω*.*

Proof. (1) We consider the diagram

$$\begin{tikzcd} {}^{\theta}\mathbf{E}_{\mathbf{W},\Omega} & {}^{\theta}\mathbf{E}'_{\Omega} \arrow[l, "p_1"'] \arrow[r, "p_2"'] & {}^{\theta}\mathbf{E}''_{\Omega} \arrow[r, "p_3"'] & {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}. \end{tikzcd}$$

Since i is a sink, we have $p_1^{-1}({}^{\theta}\mathbf{E}_{\mathbf{W},\Omega,r}) = p_2^{-1}p_3^{-1}(\mathbf{E}_{\mathbf{V},\Omega,r+1})$ for any integer r. Especially, for $L \in {}^{\theta}\mathscr{P}_{\mathbf{W},\Omega}$, $p_3p_2(p_1^{-1}\operatorname{Supp} L) \subset {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\geq\varepsilon_i(L)+1}$. For r, set ${}^{\theta}\mathbf{E}''_{\Omega,r} = p_3^{-1}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,r})$. Then $p_2^{-1}({}^{\theta}\mathbf{E}''_{\Omega,r}) = p_1^{-1}({}^{\theta}\mathbf{E}_{\mathbf{W},\Omega,r-1})$. We set ${}^{\theta}\mathbf{E}''_{\Omega,\leq r}\colon = \cup_{r' \leq r}{}^{\theta}\mathbf{E}''_{\Omega,r'}$. Then ${}^{\theta}\mathbf{E}''_{\Omega,\leq r}$ is an open subset of ${}^{\theta}\mathbf{E}''_{\Omega}$. If $p_3(x, V) = x \in {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,r}$, V_i is a one-codimensional subspace of $\mathbf{V}_i$ which contains the $(\dim \mathbf{V}_i - r)$-dimensinal subspace $\sum_{\mathrm{in}(h)=i} \operatorname{Im} x_h$ of $\mathbf{V}_i$. Therefore ${}^{\theta}\mathbf{E}''_{\Omega,r} \to {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,r}$ is a $\mathbf{P}^{r-1}$-bundle. For $L \in {}^{\theta}\mathscr{P}_{\mathbf{W},\Omega}$, there is a unique simple perverse sheaf $L'' \in \mathscr{D}({}^{\theta}\mathbf{E}''_{\Omega})$ such that $p_1^*L[d_{p_1} - d_{p_2}] = p_2^*L''$ and $(p_3)_*L'' = F_i(L)$. For $x \in {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(L)+1}$, the action of the stabilizer ${}^{\theta}G_{\mathbf{V},x} \subset {}^{\theta}\mathbf{G}_{\mathbf{V}}$ of x on $p_3^{-1}(x)$ is transitive. Since L'' is ${}^{\theta}\mathbf{G}_{\mathbf{V}}$-equivariant, L'' is a constant sheaf on any fibers of p_3 over ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(L)+1}$.
We restrict L'' to the open subset ${}^{\theta}\mathbf{E}''_{\Omega,\leq\varepsilon_i(L)+1}$. There exists a nuique simple perverse sheaf J_0 on ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L)+1}$ such that $L''|_{{}^{\theta}\mathbf{E}''_{\Omega,\leq\varepsilon_i(L)+1}} = p_3^*J_0[\varepsilon_i(L)]$. Hence $(p_3)_*L''|_{{}^{\theta}\mathbf{E}''_{\Omega,\leq\varepsilon_i(L)+1}} = (p_3)_*p_3^*J_0[\varepsilon_i(L)] = [\varepsilon_i(L)+1]_v J_0$. Let L_0 be the minimal extension of J_0. Then L_0 is a simple perverse sheaf on ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega}$. Since F_iL is semisimple, we have

$$F_i(L) = [\varepsilon_i(L)+1]_v(L_0) + \sum a_{L'}(L'),$$

where $L' \in {}^{\theta}\mathscr{P}_{\mathbf{V},\Omega}$ satisfies $\operatorname{Supp}(L') \subset {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,>\varepsilon_i(L)+1}$, or $\varepsilon_i(L') > \varepsilon_i(L)$.
To prove $a_{L'} \in v^{2-\varepsilon_i(L')}\mathbb{Z}[v]$, we restrict $\mathbf{R}\mathscr{H}om((p_3)_*L'', L')$ to the open subset ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')}$. Write $F_iL = \oplus_{J \in {}^{\theta}\mathscr{P}_{\mathbf{V},\Omega}} J \otimes M_J$, where $M_J \in \mathscr{D}(\mathrm{pt})$ is the multiplicity space of J in the expansion of F_iL. Then

$$\begin{aligned} \mathbf{R}\mathscr{H}om((p_3)_*L'', L')|_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')}} &= \bigoplus_J \mathbf{R}\mathscr{H}om(J, L')|_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')}} \otimes M_J^* \\ &\supset \mathbf{R}\mathscr{H}om(L', L')|_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')}} \otimes M_{L'}^*, \end{aligned}$$

On the other hand, since p_3 is a $\mathbf{P}^{\varepsilon_i(L')-1}$-bundle on ${}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(L')}$ and $\operatorname{Supp}(L') \cap {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')} \subset {}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(L')}$, we have

$$\begin{aligned} & \mathbf{R}\mathscr{H}om((p_3)_*L'', L')|_{{}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')}} \\ = \; & (p_3)_*\,\mathbf{R}\mathscr{H}om(L'', p_3^!L')|_{p_3^{-1}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')})} \\ = \; & (p_3)_*\,\mathbf{R}\mathscr{H}om(L'', p_3^*L'[\varepsilon_i(L')-1])|_{p_3^{-1}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')})}[\varepsilon_i(L')-1]. \end{aligned}$$

Since $p_3^*L'[\varepsilon_i(L')-1]|_{p_3^{-1}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')})}$ is a perverse sheaf, we have

$$\mathbf{R}\mathscr{H}om(L'', p^*L'[\varepsilon_i(L')-1])|_{p_3^{-1}({}^{\theta}\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')})} \in \mathscr{D}^{\geq 0}$$

by Lemma 2.16. Moreover since $\mathrm{Supp}(L'') \supsetneq \mathrm{Supp}(p_3^*L')$, we have

$$H^0(\mathbf{R}\mathscr{H}om(L'', p_3^*L'[\varepsilon_i(L')-1])|_{p_3^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')})}) = 0.$$

Therefore $\mathbf{R}\mathscr{H}om(L'', p^*L'[\varepsilon_i(L')-1])|_{p_3^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L')})} \in \mathscr{D}^{>0}(\mathrm{pt})$ and its direct image of p_3 is contained in $\mathscr{D}^{>0}$. Thus we obtain $\mathbf{R}\mathscr{H}om((p_3)_*L'', L')|_{{}^\theta\mathbf{E}_{\mathbf{V},\leq\varepsilon_i(L')}} \in \mathscr{D}^{>1-\varepsilon_i(L')}$.
Since $H^0(\mathbf{R}\mathscr{H}om(L', L')) \neq 0$, we conclude $M_{L'}^* \in \mathscr{D}^{>1-\varepsilon_i(L')}(\mathrm{pt})$. Hence $a_{L'} \in v^{2-\varepsilon_i(L')}\mathbb{Z}[v]$.
(2) Recall the following diagram:

$$\begin{tikzcd}
\mathbf{E}_{\mathbf{T},\Omega} \times {}^\theta\mathbf{E}_{\mathbf{W},\Omega} & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega \arrow[l, "p"'] \arrow[r, hook, "\iota"] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega}
\end{tikzcd}.$$

Since i is a sink, for a fixed $x_{\mathbf{W}} \in {}^\theta\mathbf{E}_{\mathbf{W},\Omega}$, $x \in {}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ is uniquely determined by the condition that U is x-stable and x induces $x_{\mathbf{W}}$ on $U/U^\perp \cong \mathbf{W}$. Therefore we have ${}^\theta\mathbf{E}_{\mathbf{W},\Omega} \cong {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega$. We have a section s of $p_1\colon {}^\theta\mathbf{E}'_\Omega \to {}^\theta\mathbf{E}_{\mathbf{W},\Omega}$ by $x_{\mathbf{W}} \mapsto (x, U, \varphi_{\mathbf{W}})$ where $\varphi_{\mathbf{W}}\colon U/U^\perp \cong \mathbf{W}$ is a given isomorphism of θ-symmetric I-graded vector spaces. We consider the following diagram:

$$\begin{tikzcd}
{}^\theta\mathbf{E}'_\Omega \arrow[r, "q\colon =p_3\circ p_2"] \arrow[d, "p_1"] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega} \arrow[r, "\sim"] \arrow[u, bend left, "s"] & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega \arrow[u, "\iota"']
\end{tikzcd}$$

For $K \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$, we have $E_iK = s^*q^*K[-\dim \mathbf{W}_i]$. Assume that $\varepsilon_i(K) > 0$. Since $\mathrm{Supp}(K) \subset {}^\theta\mathbf{E}_{\mathbf{V},\Omega,\geq\varepsilon_i(K)}$, $K|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(K)}}$ is a simple perverse sheaf. Since q is smooth on ${}^\theta\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(K)}$, the restriction $q^*K[d_q]|_{q^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(K)})}$ is a ${}^\theta\mathbf{G}_{\mathbf{V}}$-equivariant perverse sheaf, where d_q is the fiber dimension of q on ${}^\theta\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(K)}$. Note that p_1 is an affine bundle on ${}^\theta\mathbf{E}_{\mathbf{W},\Omega,\varepsilon_i(K)-1}$. If $x \in {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega$ induces $x_{\mathbf{W}} \in {}^\theta\mathbf{E}_{\mathbf{W},\Omega,\varepsilon_i(K)-1}$, the stabilizer ${}^\theta G_{\mathbf{V}}$ acts transitively on the fiber of p_1 at $x_{\mathbf{W}}$. Thus $q^*K[d_q]|_{q^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(K)})}$ is constant on any fibers of p_1. Hence $s^*q^*K[d_q - d_{p_1}]|_{{}^\theta\mathbf{E}_{\mathbf{W},\Omega,\varepsilon_i(K)-1}}$ is a simple perverse sheaf. Here

$$d_{p_1} - d_q = d_{p_1} - d_{p_2} - (\varepsilon_i(K) - 1) = \dim \mathbf{W}_i + 1 - \varepsilon_i(K).$$

Therefore we obtain

$$E_i(K) = v^{1-\varepsilon_i(K)}(K_0) + \sum_{K' \in {}^\theta\mathscr{P}_{\mathbf{W},\Omega}\colon \varepsilon_i(K') > \varepsilon_i(K)-1} b_{K'}(K'),$$

where K_0 is the minimal extension of $s^*q^*K[d_q - d_{p_1}]|_{{}^\theta\mathbf{E}_{\mathbf{W},\Omega,\varepsilon_i(K)-1}}$.
We shall prove $b_{K'} \in v^{1-\varepsilon_i(K')}\mathbb{Z}[v]$.
Since $q^*K[-\dim \mathbf{W}_i]$ and $p_1^*E_iK$ are constant along the fibers of p_1, and $s^*q^*K[-\dim \mathbf{W}_i] = s^*p_1^*E_iK$, we obtain $q^*K[-\dim \mathbf{W}_i] = p_1^*E_iK$. We have $q^*K[-\dim \mathbf{W}_i] = \oplus_{K''} p_1^*K'' \otimes M_{K''}$, where $M_{K''}$ is the multiplicity space of K'' in E_iK. Since there is a unique semisimple object $L_{K''} \in \mathscr{D}({}^\theta\mathbf{E}''_\Omega)$ such that $p_1^*K'' = p_2^*L_{K''}$, we have $p_2^*p_3^*K[-\dim \mathbf{W}_i] = \oplus_{K''} p_2^*L_{K''} \otimes M_{K''}$.
We obtain $p_3^*K[-\dim \mathbf{W}_i] = \oplus_{K''} L_{K''} \otimes M_{K''}$.

Therefore we have

$$\begin{aligned}
& \oplus_{K''}\mathbf{R}\mathrm{Hom}(L_{K''}|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}, L_{K'}|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}) \otimes M^*_{K''} \\
= \;& \mathbf{R}\mathrm{Hom}(p_3^*K[-\dim\mathbf{W}_i]|_{p_3^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1})}, L_{K'}|_{p_3^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1})}) \\
= \;& \mathbf{R}\mathrm{Hom}(K[-\dim\mathbf{W}_i]|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}, (p_3)_*L_{K'}|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}) \\
= \;& \mathbf{R}\mathrm{Hom}(K[-\dim\mathbf{W}_i]|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}, F_i(K')|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}[-\dim\mathbf{W}_i]) \\
= \;& \mathbf{R}\mathrm{Hom}(K|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(K')+1}}, F_i(K')|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}).
\end{aligned}$$

By the claim of (1), $F_iK'|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')}} = [\varepsilon_i(K')+1]_v\tilde{F}_iK'|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}} \in {}^p\mathscr{D}^{\geq-\varepsilon_i(K')}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1})$. Since $\mathrm{Supp}(K) \subsetneq \mathrm{Supp}(F_iK')$, we have $\mathbf{R}\mathrm{Hom}(K|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}, F_i(K')|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K')+1}}) \in \mathscr{D}^{\geq 1-\varepsilon_i(K')}$. Therefore $\mathbf{R}\mathrm{Hom}(L_{K'}, L_{K'})\otimes M^*_{K'} \in \mathscr{D}^{\geq 1-\varepsilon_i(K')}$, which implies $M^*_{K'} \in \mathscr{D}^{\geq 1-\varepsilon_i(K')}$.
Hence $b_{K'} \in v^{1-\varepsilon_i(K')}\mathbb{Z}[v]$ is proved.
In the case $\varepsilon_i(K) = 0$, we can prove similarly $b_{K'} \in v^{1-\varepsilon_i(K')}\mathbb{Z}[v]$. □

Lemma 5.4. *Suppose* $\mathrm{wt}\,\mathbf{V} \neq 0$. *For any* $L \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$, *there exists* $i \in I$ *such that* $\varepsilon_i(L) > 0$.

Proof. If $\mathbf{V} \neq \{0\}$, there exists an integer d, $\mathbf{i} = (i_1, \ldots, i_{2m})$ and $\mathbf{a}$ such that $L[d]$ appears in a direct summand of ${}^\theta L_{\mathbf{i},\mathbf{a};\Omega}$. We may assume $a_1 > 0$. Then, taking Ω such that i_1 is a sink, we have $\mathrm{Supp}(L) \subset \mathrm{Supp}({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) \subset {}^\theta\mathbf{E}_{\mathbf{V},\Omega,\geq 1}$. By the definition of ε_i, we have $\varepsilon_{i_1}(L) \neq 0$. □

Lemma 5.5. *For* $L \in \mathscr{P}_{\mathbf{V}}$, *we have* $\tilde{E}_i\tilde{F}_i(L) = (L)$, *and if* $\tilde{E}_i(L) \neq 0$, *we have* $\tilde{F}_i\tilde{E}_i(L) = L$.

Proof. We assume that i is a sink.
Recall the following diagram:

$$\begin{tikzcd}
{}^\theta\mathbf{E}'_\Omega \arrow[r, "q\colon = p_3\circ p_2"] \arrow[d, "p_1"] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega} \\
{}^\theta\mathbf{E}_{\mathbf{W},\Omega} \arrow[r, "\sim"] \arrow[u, bend left, "s"] & {}^\theta\mathbf{E}(\mathbf{W},\mathbf{V})_\Omega \arrow[u, "\iota"']
\end{tikzcd}$$

For $L \in {}^\theta\mathscr{P}_{\mathbf{W},\Omega}$, take simple perverse sheaf $L'' \in \mathscr{D}({}^\theta\mathbf{E}''_\Omega)$ such that $p_1^*L[\dim\mathbf{W}_i] = p_2^*L''$ and $(p_3)_!L'' = F_iL$, then $(p_3)_!L'' \cong [\varepsilon_i(L)+1]_v\tilde{F}_iL$ on ${}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L)+1}$. On the other hand, since $L'' \cong p_3^*\tilde{F}_iL[\varepsilon_i(L)]$ on $p_3^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(L)+1})$, we have $q^*\tilde{F}_iL \cong p_2^*L''[-\varepsilon_i(L)] = p_1^*L[\dim\mathbf{W}_i - \varepsilon_i(L)]$ on $p_1^{-1}({}^\theta\mathbf{E}_{\mathbf{W},\Omega,\leq\varepsilon_i(L)})$. Then we have $s^*q^*\tilde{F}_iL = L[\dim\mathbf{W}_i - \varepsilon_i(L)]$ on ${}^\theta\mathbf{E}_{\mathbf{W},\Omega,\leq\varepsilon_i(L)+1}$. Note that $E_i\tilde{F}_iL = s^*q^*\tilde{F}_iL[-\dim\mathbf{W}_i]$. We obtain $E_i(\tilde{F}_iL) = L[-\varepsilon_i(L)]$ on ${}^\theta\mathbf{E}_{\mathbf{W},\Omega,\leq\varepsilon_i(L)}$. Hence $\tilde{E}_i\tilde{F}_i(L) = (L)$.
Conversely, take $K \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$ such that $\varepsilon_i(K) > 0$. By the similar argument in the proof of Theorem 5.3(2), we have $p_1^*E_iK = q^*K[-\dim\mathbf{W}_i]$. Hence we obtain $p_1^*\tilde{E}_iK[\dim\mathbf{W}_i] = q^*K[\varepsilon_i(K)-1]$ on $p_1^{-1}({}^\theta\mathbf{E}_{\mathbf{W},\Omega,\varepsilon_i(K)-1})$. Since $p_3^*K[\varepsilon_i(K)-1]$ is a simple perverse sheaf on $p_3^{-1}({}^\theta\mathbf{E}_{\mathbf{V},\Omega,\varepsilon_i(K)})$, we have $F_i\tilde{E}_iK = (p_3)_*p_3^*K[\varepsilon_i(K)-1] = [\varepsilon_i(K)]_vK$ on ${}^\theta\mathbf{E}_{\mathbf{V},\Omega,\leq\varepsilon_i(K)}$. Then we have $\tilde{F}_i\tilde{E}_i(K) = (K)$. □

5.4. **Verdier duality functor.** The Verdier duality functor $D\colon \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega}) \to \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$ satisfies $D(L[d]) = D(L)[-d]$ for $L \in \mathscr{D}({}^\theta\mathbf{E}_{\mathbf{V},\Omega})$, $d \in \mathbb{Z}$. Then D induces the involution $v^{\pm 1} \mapsto v^{\mp 1}$.

Proposition 5.6.

(i) $D({}^\theta L_{\mathbf{i},\mathbf{a};\Omega}) = {}^\theta L_{\mathbf{i},\mathbf{a};\Omega}[2\dim {}^\theta\widetilde{\mathcal{F}}_{\mathbf{i},\mathbf{a};\Omega}]$.

(ii) For any $L \in {}^\theta\mathscr{Q}_{\mathbf{V},\Omega}$, *we have* $D(F_iL) = F_iD(L)$.

(iii) For any $L \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$, *we have* $D(L) \cong L$.

Proof. (i) and (ii) follow from the general property of the Verdier duality functor (see Lemma 2.15).

To prove (iii), we use the induction on $\mathrm{wt}\,\mathbf{V}$.

When $\mathrm{wt}\,\mathbf{V} = 0$, the claim is clear by ${}^\theta\mathscr{P}_{\mathbf{V},\Omega} = \{\mathbf{1}_{\mathrm{pt}}\}$ and $D(\mathbf{1}_{\mathrm{pt}}) = \mathbf{1}_{\mathrm{pt}}$.

Suppose $\mathrm{wt}\,\mathbf{V} \neq 0$. By Lemma 5.4, there exists i such that $\varepsilon_i(L) > 0$. We shall prove $D(L) = L$ by the descending induction on $\varepsilon_i(L)$. By Theorem 5.3 and Lemma 5.5, we have

$$F_i(\tilde{E}_iL) = [\varepsilon_i(L)]_v(L) + \sum_{L' \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}:\, \varepsilon_i(L') > \varepsilon_i(L)} a_{L'}(L').$$

By the induction hypothesis on $\mathrm{wt}\,\mathbf{V}$, $D(\tilde{E}_iL) = \tilde{E}_iL$. Hence the lefthand side is D-invariant by (ii). We restrict $F_i(\tilde{E}_iL)$ on the open subset ${}^\theta\mathbf{E}_{\mathbf{V},\Omega,\le\varepsilon_i(L)}$. Then it is isomorphic to $[\varepsilon_iL]_v(L)|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\le\varepsilon_i(L)}}$ and D-invariant. Since L is the minimal extension of $L|_{{}^\theta\mathbf{E}_{\mathbf{V},\Omega,\le\varepsilon_i(L)}}$, L is D-invariant. ☐

Remark 5.7. By the result of (iii), we have $a_{L'}(v) = a_{L'}(v^{-1})$ in Theorem 5.3 (1).

Lemma 5.8. *For* $L \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$, *we have*

$$F_i^{(a)}(L) = \begin{bmatrix} \varepsilon_i(L)+a \\ a \end{bmatrix}_v (\tilde{F}_i^aL) + \sum_{L' :\, \varepsilon_i(L') > \varepsilon_i(L)+a} c_{L'}(L')$$

with $c_{L'} \in \mathbb{Z}[v, v^{-1}]$.

Proof. We shall prove the claim by the induction on a. If $a = 1$, the claim follows from Theorem 5.3. If $a > 1$, by the induction hypothesis and Theorem 5.3, we have

$$\begin{aligned} F_iF_i^{(a)}(L) &= \begin{bmatrix} \varepsilon_i(L)+a \\ a \end{bmatrix}_v F_i(\tilde{F}_i^aL) + \sum_{L' :\, \varepsilon_i(L') > \varepsilon_i(L)+a} c_{L'}F_i(L') \\ &= [a+1]_v \left(\begin{bmatrix} \varepsilon_i(L)+a+1 \\ a+1 \end{bmatrix}_v (\tilde{F}_i^{a+1}L) + \sum_{L'' :\, \varepsilon_i(L'') > \varepsilon_i(L)+a+1} d_{L''}(L'') \right), \end{aligned}$$

where $d_{L''} \in \mathbb{Q}(v)$. Hence

$$F_i^{(a+1)}L = \begin{bmatrix} \varepsilon_i(L)+a+1 \\ a+1 \end{bmatrix}_v (\tilde{F}_i^{a+1}L) + \sum_{L'' :\, \varepsilon_i(L'') > \varepsilon_i(L)+a+1} d_{L''}(L'').$$

On the other hand, since $F_i^{(a+1)}L = \mathbf{1}_{\mathbf{S}_i^{a+1}} * L[d_{a+1}]$ is semisimple, we conclude $d_{L''} \in \mathbb{Z}[v, v^{-1}]$. ☐

Proposition 5.9. *We have* ${}^\theta K = \sum \mathbb{Z}[v, v^{-1}]F_{i_1}^{(a_1)} \cdots F_{i_k}^{(a_k)}\mathbf{1}_{\{\mathrm{pt}\}}$.

Proof. For $L \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}$ such that $\mathrm{wt}\,\mathbf{V} \neq 0$, there exists i such that $\varepsilon_i(L) > 0$. We shall prove that (L) is contained in $\sum \mathbb{Z}[v, v^{-1}]F_{i_1}^{(a_1)} \cdots F_{i_k}^{(a_k)}\mathbf{1}_{\{\mathrm{pt}\}}$ by the induction on $\mathrm{wt}\,\mathbf{V}$ and the descending induction on $\varepsilon_i(L)$. We have

$$F_i^{(\varepsilon_i(L))}(\tilde{E}_i^{\varepsilon_i(L)}L) = (L) + \sum_{L' \in {}^\theta\mathscr{P}_{\mathbf{V},\Omega}:\, \varepsilon_i(L') > \varepsilon_i(L)} c_{L'}(L')$$

by Lemma 5.8 and Lemma 5.5. By the induction hypothesis, we have $c_{L'}(L')$ and $\tilde{E}_i^{\varepsilon_i(L)}L$ are contained in $\sum \mathbb{Z}[v,v^{-1}]F_{i_1}^{(a_1)}\cdots F_{i_k}^{(a_k)}\mathbf{1}_{\{\mathrm{pt}\}}$. Thus $(L) \in \sum \mathbb{Z}[v,v^{-1}]F_{i_1}^{(a_1)}\cdots F_{i_k}^{(a_k)}\mathbf{1}_{\{\mathrm{pt}\}}$. □

5.5. **Main Theorem.** Let us recall

$$^{\theta}K' := \sum_{(\mathbf{i},\mathbf{a})} \mathbb{Z}[v,v^{-1}](^{\theta}L_{\mathbf{i},\mathbf{a};\Omega}) = \sum \mathbb{Z}[v,v^{-1}]F_{i_1}^{(a_1)}\cdots F_{i_k}^{(a_k)}\mathbf{1}_{\{\mathrm{pt}\}} \subset {}^{\theta}K.$$

Theorem 5.10.

(i) $^{\theta}K = {}^{\theta}K'$.

(ii) For $L \in {}^{\theta}\mathscr{P}_{\mathbf{V}}$, *we define* $\mathrm{wt}(L) = -\mathrm{wt}\,\mathbf{V}$. *Then* $(\mathrm{wt}, \tilde{E}_i, \tilde{F}_i, \varepsilon_i)$ *gives a crystal structure on* $^{\theta}\mathscr{P} := \sqcup_{\mathbf{V}} {}^{\theta}\mathscr{P}_{\mathbf{V}}$ *in the sence of section 2.3. Here* $\mathbf{V}$ *runs over all isomorphism classes of* θ*-symmetric* I*-graded vector spaces.*

(iii) Let $\mathcal{L}$ *be the* $\mathbf{A}_0$*-submodule* $\sum_{(L)\in^{\theta}\mathscr{P}} \mathbf{A}_0(L)$ *of* $^{\theta}K$. *Then* $\{(L) \mod v\mathcal{L} | L \in {}^{\theta}\mathscr{P}\}$ *gives a crystal basis of* $^{\theta}K$. *Especially, the actions of modified root operators* $\tilde{E}_i$ *and* $\tilde{F}_i$ *on* $\mathcal{L}/v\mathcal{L}$ *are compatible with the actions of* $\tilde{E}_i$ *and* $\tilde{F}_i$ *on* $^{\theta}\mathscr{P}$ *introduced in Theorem 5.3.*

Proof. (i) is nothing but Proposition 5.9.
(ii) By the definition of $\varepsilon_i(L), \tilde{F}_i$ and $\tilde{E}_i$, and Lemma 5.5, we conculde that $(\mathrm{wt}, \tilde{E}_i, \tilde{F}_i, \varepsilon_i)$ gives a crystal structure on $^{\theta}\mathscr{P} := \sqcup_{\mathbf{V}} {}^{\theta}\mathscr{P}_{\mathbf{V}}$ in the sence of section 2.3(i)-(iv). By the estimates in Theorem 5.3, the actions of E_i and F_i on (L) $(L \in {}^{\theta}\mathscr{P})$ satisfy the conditions 2.1-2.6 in section 2.3. Thus we obtain the claim.
(iii) follows from Theorem 2.14. □

Lemma 5.11. *We have* $\{v \in {}^{\theta}K \mid E_i v = 0$ *for any* $i \in I\} = \mathbb{Z}[v,v^{-1}]\mathbf{1}_{\{\mathrm{pt}\}}$.

Proof. Suppose that $E_i\left(\sum a_L(L)\right) = 0$ for any L. Then $a_L \in v^c\mathbb{Z}[v]$ for some c. Put $\widetilde{a_L} = v^{-c}a_L \in \mathbb{Z}[v]$. By the definition of the modified root operators and Theorem 5.10(iii), we have $\tilde{E}_i\left(\sum \tilde{a_L}(L)\right) = 0$. Specializing v to 0, we have $\widetilde{a_L}(0) = 0$ if $\tilde{E}_i L \neq 0$. But for any L such that $\mathrm{wt}(L) \neq 0$, there exists $i \in I$ such that $\varepsilon_i(L) > 0$. Hence we obtain $\widetilde{a_L} \in v\mathbb{Z}[v]$ and hence $a_L \in v^{c+1}\mathbb{Z}[v]$. By the induction on c, we have $a_L \in v^c\mathbb{Z}[v]$ for any c. Thus we conclude $a_L = 0$ for $\mathrm{wt}(L) \neq 0$. □

Theorem 5.12.

(i) $^{\theta}K \otimes_{\mathbb{Z}[v,v^{-1}]} \mathbb{Q}(v) \cong V_\theta(0)$ *as a* $B_\theta(\mathfrak{g})$*-module. The involution induced by the Verdier duality functor coincides with the bar involution on* $V_\theta(0)$.

(ii) $\{(L) \mid L \in {}^{\theta}\mathscr{P}\}$ *gives the lower global basis on* $V_\theta(0)$.

Proof. (i) By Proposition 5.1, to check the defining relations of $B_\theta(\mathfrak{g})$, we only need to prove the v-Serre relations. Put

$$S_e = \sum_{k=0}^{b}(-1)^k E_i^{(k)}E_jE_i^{(b-k)}, \quad S_f = \sum_{k=0}^{b}(-1)^k F_i^{(k)}F_jF_i^{(b-k)}$$

and note that $F_kS_e = S_eF_k$ and $E_kS_f = S_fE_k$ for any $k \in I$.
Since $^{\theta}K_\Omega$ is generated by $F_k^{(n)}$'s from $\phi := \mathbf{1}_{\{\mathrm{pt}\}}$ and $S_e\phi = 0$, we have $S_ev = 0$ for any $v \in {}^{\theta}K_\Omega$. We show $S_f(L) = 0$ for any $L \in {}^{\theta}\mathscr{P}_{\mathbf{V},\Omega}$ by the induction on $\mathrm{wt}\,\mathbf{V}$. If $\mathrm{wt}(S_f(L)) \neq 0$, we have we have $E_kS_f(L) = S_fE_k(L) = 0$ for any $k \in I$ by applying the induction hypothesis to $E_k(L)$. Since $\mathrm{wt}(S_f(L)) \neq 0$, we have $S_f(L) = 0$ by Lemma 5.11. Hence $^{\theta}K$ is a $B_\theta(\mathfrak{g})$-module. Note that $T_i\mathbf{1}_{\{\mathrm{pt}\}} = \mathbf{1}_{\{\mathrm{pt}\}}$ for any $i \in I$. We conclude $^{\theta}K \cong V_\theta(0)$ by Lemma 5.11 and the characterization of $V_\theta(0)$ in Proposition 2.10.

(ii) We already know that $\mathcal{L} = \sum_{L \in {}^\theta\mathscr{P}} \mathbf{A}_0(L)$ is a crystal lattice and $\{(L) \mod v\mathcal{L}\}$ is a basis of $\mathcal{L}/v\mathcal{L}$. Note that $\sum_{L \in {}^\theta\mathscr{P}} \mathbb{Z}[v, v^{-1}](L)$ is stable under the actions of E_i's and $F_i^{(a)}$'s by Lemma 5.8 and L is D-invariant, namely bar-invariant. Moreover $\{(L) \mid L \in {}^\theta\mathscr{P}\}$ is a basis of the $\mathbf{A}_0$-module $\mathcal{L}$ and also a basis of the $\mathbb{Z}[v, v^{-1}]$-module ${}^\theta K$. Hence we conclude that $\{(L) \mid L \in {}^\theta\mathscr{P}\}$ gives the lower global basis on $V_\theta(0)$. ☐

Corollary 5.13. *For any Kac-Moody algebra $\mathfrak{g}$ with a symmetric Cartan matrix, the $B_\theta(\mathfrak{g})$-module $V_\theta(0)$ has a crystal basis and a lower global basis, namely Conjecture 2.11 and Conjecture 2.12 is true if $\lambda = 0$.*

Example 5.14. Let us consider the case $\mathfrak{g} = \mathfrak{sl}_3$, $I = \{\pm 1\}$ and $\theta(i) = -i$. Fix a θ-symmetric orientation $-1 \xrightarrow{\Omega} 1$. For a θ-symmetric I-graded vector space $\mathbf{V}$ such that $\mathrm{wt}(\mathbf{V}) = n(\alpha_{-1} + \alpha_1)$, ${}^\theta\mathbf{E}_{\mathbf{V},\Omega}$ is the set of skew symmetric matrix x of size n. Its ${}^\theta\mathbf{G}_{\mathbf{V}}$-orbits are parametrized by the rank $2r$ ($0 \le r \le \lfloor \frac{n}{2} \rfloor$) of x. We denote $\mathcal{O}_r^n$ by the orbit consisting of $n \times n$ skew symmetric matrices x of rank $2r$. Note that any simple local system on each ${}^\theta\mathbf{G}_{\mathbf{V}}$-orbit is trivial. Let us denote IC_r^n by the simple perverse sheaves corresponding to the orbit $\mathcal{O}_r^n$. Note that $\varepsilon_1(\mathrm{IC}_r^n) = n - 2r$.

Let $\mathbf{W}$ be a θ-symmetric I-graded vector space such that $\mathrm{wt}(\mathbf{W}) = (n-1)(\alpha_{-1} + \alpha_1)$. We consider the diagram:

$$\begin{tikzcd}
{}^\theta\mathbf{E}_{\mathbf{W},\Omega} & {}^\theta\mathbf{E}'_\Omega \arrow[l, "p_1"'] \arrow[r, "p_2"'] & {}^\theta\mathbf{E}''_\Omega \arrow[r, "p_3"'] & {}^\theta\mathbf{E}_{\mathbf{V},\Omega}.
\end{tikzcd}$$

Note that the fibers of p_3 on $\mathcal{O}_r^n$ is isomorphic to $\mathbf{P}^{n-1-2r}$. Then

$$F_1(\mathrm{IC}_r^{n-1}) = [n-2r]_v(\mathrm{IC}_r^n) + \sum_{k=0}^{r-1} a_{k,n}(\mathrm{IC}_k^n)$$

where $a_{k,n} \in v^{2-n+2k}\mathbb{Z}[v]$. We obtain the crystal graph:

$$\begin{tikzcd}
 & & & & \mathrm{IC}_0^4 \arrow[r, shift left, "1"] \arrow[r, shift right, "-1"'] & \mathrm{IC}_0^5 \cdots \\
 & & \mathrm{IC}_0^2 \arrow[r, shift left, "1"] \arrow[r, shift right, "-1"'] & \mathrm{IC}_0^3 \arrow[ur, "1"] \arrow[dr, "-1"] & & \\
\mathrm{IC}_0^0 \arrow[r, shift left, "1"] \arrow[r, shift right, "-1"'] & \mathrm{IC}_0^1 \arrow[ur, "1"] \arrow[dr, "-1"'] & & & \mathrm{IC}_1^4 \arrow[r, shift left, "1"] \arrow[r, shift right, "-1"'] & \mathrm{IC}_1^5 \cdots \\
 & & \mathrm{IC}_1^2 \arrow[r, shift left, "1"] \arrow[r, shift right, "-1"'] & \mathrm{IC}_1^3 \arrow[ur, "-1"] \arrow[dr, "-1"'] & & \\
 & & & & \mathrm{IC}_2^4 \arrow[r, shift left, "1"] \arrow[r, shift right, "-1"'] & \mathrm{IC}_2^5 \cdots
\end{tikzcd}$$

Therefore we recover the crystal graph parametrized by "θ-restricted multi-segments" in [EK2, Example 4.7 (1)].

REFERENCES

[Ari] Susumu Ariki, *On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$*, J. Math. Kyoto Univ. 36 (1996), no. 4, 789–808.

[BBD] Aleksandr A. Beilinson, Joseph N. Bernstein and Pierre Deligne, *Faisceaux pervers*, Astérisque, **100**, Soc. Math. France, Paris, 1982

[EK1] Naoya Enomoto and Masaki Kashiwara, *Symmetric crystals and the affine Hecke algebras of type B*, Proc. Japan. Acad. (2006), 82, no.8, 131-136.

[EK2] ———, *Symmetric Crystals for $\mathfrak{gl}_\infty$*, to appear in Publ. RIMS. (2008), vol.44, no.3, 837–891, arxiv:math.QA/0704.2817.

[EK3] ———, *Symmetric Crystals and LLT-Ariki Type Conjectures for the Affine Hecke Algebras of Type B*, RIMS kokyuroku bessatsu (2008), B8, 1–20, arxiv:math.RT/0705.3938

[Ka1] Masaki Kashiwara, *On crystal bases of the q-analogue of universal enveloping algebras*, Duke Math. J. 63 (1991), no. 2, 465–516.

[Ka2] ———, *Global crystal bases of quantum groups*, Duke Math. J. **69**(1993), no.2, 455-485

[KS] Masaki Kashiwara and Pierre Schapira, *Sheaves on manifolds*, Grundlehren der mathematischen Wissenschaften, vol.292, Springer, 1990

[Kt] Syu Kato, *An exotic Deligne-Langlands correspondence for symplectic groups*, arxiv:math.RT/0601155.

[Lau] Gérard Laumon, *Transformation de Fourier, constantes d'équations fonctionnelles et conjecture de Weil*, Publ. Math. I.H.E.S, 65 (1987), 131–210

[Lus1] George Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc., 3 (1990), no. 2, 447–498.

[Lus2] ———, *Quivers, Perverse sheaves and Quantized enveloping algebras*, J. Amer. Math. Soc., 4-2(1991), 365–421

[Lus3] ———, *Introduction to Quantum Groups*, Birkhäuser, 1993

[Lus4] ———, *Study of Perverse Sheaves Arising from Graded Lie Algebras*, Adv. Math., **112** (1995), 147–217

[Lus5] ———, *Graded Lie Algebras and Intersection Cohomology*, arxiv:math.RT/0604535.

[LLT] Alain Lascoux, Bernard Leclerc and Jean Y. Thibon, *Hecke algebras at roots of unity and crystal bases of quantum affine algebras*, Comm. Math. Phys. 181 (1996), no. 1, 205–263.

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606–8502, Japan

E-mail address: `henon@kurims.kyoto-u.ac.jp`